



AGREEMENT ON THE CONSERVATION OF
AFRICAN-EURASIAN MIGRATORY WATERBIRDS



5th SESSION OF THE MEETING OF THE PARTIES
14 – 18 May 2012, La Rochelle, France

“Migratory waterbirds and people - sharing wetlands”

RESOLUTION 5.10

REVISION AND ADOPTION OF CONSERVATION GUIDELINES

Recalling Article IV paragraph 4 of the Agreement, and paragraph 7.3 of the Agreement’s Action Plan, which require the development and review of conservation guidelines in order to assist Contracting Parties with their implementation of the Agreement,

Further recalling Resolutions 1.10, 2.3 and 4.13, which adopted 13 conservation guidelines focusing on various aspects of waterbird conservation practice,

Noting that these conservation guidelines, although legally non-binding, provide a common framework for action, which aids the coherent implementation of the Agreement by the Contracting Parties to the Agreement, as well as other Range States and interested parties,

Noting, however, that in the 17 years since the finalisation of the Agreement text, the internet has enabled rapid access to a wide range of relevant information and guidance, notwithstanding problems of internet access in some areas, *further noting* that a review by the Ramsar Convention of uses of its technical guidance showed widespread awareness amongst Contracting Parties and other stakeholders and *recognising* that AEWA’s Conservation Guidelines are an effective means of disseminating information to Contracting Parties and other stakeholders,

Recognising the work of the UNEP/AEWA Secretariat and the Technical Committee over the past quadrennium to review previously adopted conservation guidelines and develop additional ones, also in collaboration with the Convention on Migratory Species (CMS) and the CMS MoU on the Conservation of African-Eurasian Birds of Prey,

Acknowledging the voluntary contribution by the Government of Switzerland and the funds provided by RWE Rhein-Ruhr Netzservice GmbH, which enabled the development of additional guidelines,

Recalling Resolution 4.3 which requested the Technical Committee to review the periods during which huntable bird populations of conservation concern covered by the Agreement return to their breeding grounds and, if needed, to provide further guidance on the implementation of paragraph 2.1.2 (a) AEWA Action Plan,

Recognising the work of the Technical Committee over the past quadriennium towards developing guidance on the definition of periods of breeding and pre-nuptial migration for migratory African-Eurasian waterbirds (document AEWA/MOP Inf. 5.3).

The Meeting of the Parties:

1. *Adopts* the following newly developed conservation guidelines:

a) Guidelines on how to avoid or mitigate impact of electricity power grids on migratory birds in the African-Eurasian region (document AEWA/MOP 5.37), and

b) Guidelines for the translocation of waterbirds for conservation purposes: complementing the IUCN guidelines (document AEWA/MOP 5.36),

in accordance with Article IV.4 of the Agreement, as guidance for the Contracting Parties in the implementation of the Agreement and its Action Plan;

2. *Adopts* the revised versions of the following previously adopted conservation guidelines:

a) Guidelines on identifying and tackling emergency situations for migratory waterbirds (document AEWA/MOP 5.35),

b) Guidelines on regulating trade in migratory waterbirds (document AEWA/MOP 5.33), and

c) Guidelines on avoidance of introductions of non-native waterbird species (document AEWA/MOP 5.34 Rev. 1);

3. *Calls upon* Contracting Parties to utilise these guidelines in a practical way that leads to a minimum of additional bureaucracy and that recognises the different social, economic and environmental conditions within the Agreement area;

4. *Instructs* the Secretariat to disseminate these guidelines to all Range States, and relevant international governmental and non-governmental organisations, and to monitor their use to the extent that this is possible;

5. *Encourages* Contracting Parties to utilise the guidance on defining periods of breeding and pre-nuptial migration for migratory African-Eurasian waterbirds with respect to the implementation of paragraph 2.1.2 (a) AEWA Action Plan as annexed to this Resolution;

6. *Requests* the Technical Committee and the Secretariat to integrate this guidance in a future revised version of the Conservation Guidelines on sustainable harvest of migratory waterbirds;

7. *Requests* the Technical Committee, as a matter of priority and in the first part of the next triennium, to undertake a critical review of the style and format of AEWA's Conservation Guidelines, *inter alia* considering the following existing issues:

7.1 the merits or otherwise of shorter information notes that might be easier to translate into local languages;

7.2 the need to target different styles or types of guidance to different audiences (e.g. government policy makers, wetland managers, other relevant stakeholders or user groups);

7.3 the merits or otherwise of regionally specific guidance;

7.4 knowledge of the extent of use of the existing guidelines and implications for the dissemination of guidance; and

7.5 the potential value of a 'guidance to guidance' format as has been developed by the Ramsar Convention's Scientific and Technical Review Panel;

8. *Further requests* the Technical Committee to make recommendations to the Standing Committee on the basis of the review described in paragraph 7 above, prior to developing further guidance in the current format for consideration by the Sixth Meeting of the Parties;

9. *Requests* the Technical Committee to continue its work to provide relevant information on African species, as mentioned in Appendix 1 of this Resolution, as knowledge of these improves through the implementation of the Action Plan for Africa (Resolution 5.9) and bring elaborated guidance to a future session of the Meeting of the Parties.

APPENDIX 1

DEFINING PERIODS OF BREEDING AND PRE-NUPTIAL MIGRATION FOR MIGRATORY AFRICAN-EURASIAN WATERBIRDS

Introduction

Section 2.1.2 of the AEWa Action Plan states:

“2.1.2 Parties with populations listed in Table 1 shall regulate the taking of birds and eggs of all populations listed in column B of Table 1. The object of such legal measures shall be to maintain or contribute to the restoration of those populations to a favourable conservation status and to ensure, on the basis of the best available knowledge of population dynamics, that any taking or other use is sustainable. Such legal measures, subject to paragraph 2.1.3 below, shall in particular:

- (a) prohibit the taking of birds belonging to the populations concerned during their various stages of reproduction and rearing and during their return to their breeding grounds if the taking has an unfavourable impact on the conservation status of the population concerned; ... “

In Resolution 4.3, the Meeting of the Parties requested the Technical Committee, *inter alia*:

5. To review the periods during which huntable bird populations of conservation concern covered by the Agreement return to their breeding grounds and, if needed, to provide further guidance on the implementation of paragraph 2.1.2 (a) AEWa Action Plan.

Previous relevant guidance

Article 7(4) of the EU Directive on the conservation of wild birds contains similar provisions:

4. Member States shall ensure that the practice of hunting, including falconry if practised, as carried on in accordance with the national measures in force, complies with the principles of wise use and ecologically balanced control of the species of birds concerned and that this practice is compatible as regards the population of these species, in particular migratory species, with the measures resulting from Article 2. **They shall see in particular that the species to which hunting laws apply are not hunted during the rearing season nor during the various stages of reproduction. In the case of migratory species, they shall see in particular that the species to which hunting regulations apply are not hunted during their period of reproduction or during their return to their rearing grounds.** Member States shall send the Commission all relevant information on the practical application of their hunting regulations. [Emphasis added]

The European Commission, with the Member States have developed guidance on defining periods of breeding and pre-nuptial migration, initially for the EU15 and more recently for all 27 Member States (European Commission 2009). That analysis used the following definition:


“Period of reproduction”

‘**Breeding season**’¹ was defined using the definition of Cramp & Simmons (1977): “*the breeding season is the period during which a species lays and incubates its eggs and rears its young to the flying stage.*” However, the ‘**reproduction period**’ not only covers the breeding season but also includes the occupation of the breeding areas as well as the period of dependence of young birds after leaving the nest (previously recognised in the 1993 Commission report on the application of the Birds Directive²).

“The following scheme, which deals with the different stages of reproduction, was agreed as an appropriate general scheme for the period of reproduction. The sequence and importance of the elements of this general scheme may vary by species according to differences in breeding biology.”

¹ This term is considered equal and better English than the term 'rearing season' used in Article 7(4).

² COM (93) 572 final. *Second report on the application of Council Directive 79/409/EEC on the conservation of wild birds.* Brussels, 24 November 1993.



“Criteria used to identify the beginning and end of the period of reproduction”

“In general, for migratory species, the stage of reproduction identifying the start of the period of reproduction is the 'occupation of the breeding sites'. However, the occupation of the breeding sites is generally difficult to use where the species is mainly locally resident or where there is a mixing of locally resident and migratory birds. In these cases, the stage identifying the start of the period of reproduction is the 'construction of the nest'. In those situations where the stage retained is difficult to recognise in the field, a mention is made to the corresponding number of decades counted from the start of egg laying (generally well known for most species).

- “In general, the stage retained to identify the end of the period of reproduction is the 'full flight of young birds', i.e. fledging of all broods including second or third broods for some species (e.g. rails / Rallidae, pigeons / Columbidae, thrushes / Turdidae). Full flight means that young birds are capable of sustained, continuous flight to a similar capacity as adult birds and corresponds to the 'independence of young birds. Nonetheless, for certain species (e.g. crows / Corvidae) the full flight occurs before 'independence of young birds'. Young birds are independent when the loss of parental care and/or feeding does not significantly lower survival prospects of young. In those situations where the 'full flight/independence of young' is difficult to establish in the field, a mention is made to the corresponding number of decades counted from the end of hatching.”

Migration

The EU adopted the following guidance with respect to determining the timing of migration (or ‘return to the breeding areas’):

“Return to the breeding areas”³

“Return to the breeding areas is an annual displacement, in one of more stages, of birds from their wintering areas back to nesting grounds. The wintering period ends with departure from the wintering areas where migrant birds have been more or less stationary since the end of the post-nuptial (autumn) migration. The return to the breeding areas is commonly called ‘pre-nuptial migration’ or ‘spring migration’.

In Europe, return migration movements are mostly directed north, northeast or northwest. This means that migrants from African winter quarters first cross the Mediterranean, then pass through central Europe on their way to their Northern European breeding areas. This migration normally takes several weeks (including breaks at resting places on the way) but individual birds can complete the journey in one or a few days. The start, end and length of the migration season in a particular country are determined by a number of biological, geographical and methodological factors.

Regarding the beginning of the pre-nuptial migration, all individuals of a species within a same region do not end their wintering period at the same time. Not only are there individual differences, but within a single wintering area, birds of different populations having different annual cycles come together. Birds belonging to northern populations, for example, often start their return flight much later than birds breeding more to the south. An extreme case is the so-called ‘leapfrog’ migration (e.g. in the Redshank): birds breeding in more northern latitudes travel greater distances and move to more southerly wintering areas than those that nest farther south.

The fact that birds leave a wintering area does not necessarily mean that they start their return migration. They can move to other wintering quarters because of changes in the local ecological conditions, exhaustion of food resources, disturbance or changes in climatic conditions. When migratory and sedentary birds of the same species coexist on the same wintering grounds, the situation can be even more complex. Thus, apparent discrepancies may arise among the data for large countries. Major differences between neighbouring regions can reflect ecological differences more than actual differences in migration timing. For example, although the southern parts of Spain (Andalucía) and Italy (Sicilia) are situated on the same latitude (37th) this does not necessarily imply similar arrival dates of migrants because different populations might be involved.

The length of the migration period does not only depend on the north-south extension of the country concerned but also on the availability and the use of resting places. A typical example concerns the Bar-tailed Godwits, which migrate from the African winter quarters to Siberian breeding areas. After a continuous flight from the Banc d'Arguin in Western Africa, they stay several weeks in the Wadden Sea. The migration period length is also determined by the quantity and the geographical range of the birds involved: a small population can pass in a few days while a numerous species with an extensive breeding range can have a prolonged migratory season encompassing several months. Moreover, the migration period can also be extended if a country is passed over by several populations with different time schedules.

Methodological reasons can also account for a short period: the start and end dates of migration are not recorded accurately because it only involves small numbers of birds which are often not noted if few observations are available (low chance of recording). As said before, availability of data differs very much from species to species (behavioural differences) and from country to country (e.g. numbers of observers).

In general, the beginning of the return migration can only be estimated by comparison of data from many different regions of the European Union, analysis of ring recoveries and consideration of arrival dates in the breeding areas.

Information defining the timing of pre-nuptial migration was based on statistics relating to populations rather than individual birds.”

³ "return to breeding areas" is taken as a synonym of "return to the rearing grounds"

“Presentation of data [on migration and breeding periods]”

“To avoid spurious precision and to allow for normal between-year variation in timing of migration and breeding events, the data presentations summarise the data on reproduction and return migration in ‘decades’ or ten-day periods (i.e. 1-10, 11-20, 21 up to 31 in each month).

A number of general principles were adopted in the gathering of data:

- Where there is a range in timing of pre-nuptial migration or breeding (as will occur in most countries of significant size), the data used relate to the earliest periods in each of the Member States concerned. This is generally relating to the southernmost parts or lowest altitudes. Likewise, for the end of the reproduction, the data used refers to the latest dates. This means that regional differences may exist for prenuptial migration and reproduction periods within the territory of one Member State, which may be relevant. ...
- Where significant between-year variation occurs on a regular basis, data from the earliest periods have always been taken.
- Where different populations of the same species migrate through a country at different times, information relating to the earliest migrating population has been used. In some cases, where different populations (i.e. different subspecies or different flyways) are clearly distinguishable in the field, their correspondent timings where given.
- Extreme, outlying and erratic data have been excluded due to their unpredictable nature and falling outside normal patterns of variation between and within years.”

Beyond Europe

In Africa, the definition of breeding seasons and migration periods become increasingly complex as the reproductive cycles of birds relate to different forms of seasonality and environmental predictability. These issues are explored in detail by Dodman & Diagona (2006) who highlighted that, in Africa, there are a range of definitional and other complexities that make the concept of simple latitudinal migration patterns of northern or temperate Eurasia difficult to apply. Such issues include multiple migratory behavioural types, such as:

- Local movers/short distance migrants;
- Rains migrants/ arid zone migrants;
- Nutrition migrants/post-roost dispersers;
- Post-breeding dispersers;
- Nomads;
- Altitudinal migrants; and
- Environmental response migrants.

Further, Dodman & Diagona (2006) stressed that whilst migration in northern climates is strongly determined by seasonal patterns, in tropical regions there are multiple different triggers for waterbird movements, including:

- Sudden availability of productive wetlands;
- Rising water levels/flooding;
- Falling water levels/edge effects; and
- Lack of rain/increasing aridity.

Dodman & Diagona (2006) further highlighted a range of practical problems in relation to defining intra-African migration:

- Many African flyways are diffuse, and not easy to specify.
- Some sites are only important irregularly, e.g. once every few years, especially temporary wetlands.

- Site networks are not always obvious, and may include large numbers of small wetlands or sites that are not used regularly.
- Several species exploit wetlands at different periods and for different reasons, such that sites cannot be maintained in a constant state; rather it is important to permit natural flooding and other cycles.
- Many waterbirds are nomadic and are not faithful to specific routes or annual seasons.
- It is difficult to monitor intra-African migrants: current procedures under the AfWC focused on coordinated biannual censuses are not effective enough in identifying migratory strategies.
- On a practical level, there are low resources and capacity for conserving intra-African migrants, whilst other issues also influence monitoring, such as inaccessibility and security.

They concluded that “the high diversity of “movement strategies” of African waterbirds and the often limited ability to predict movements render their management and conservation quite difficult. The life cycles and movements of most African waterbirds are not precisely known, and the networks of key sites not well determined.

Recommendations related to section 2 of AEWA’s Action Plan

The Technical Committee recommends that for the purposes of the implementation of the obligations of section 2 of the AEWA Action Plan, one of the following options are pursued as appropriate:

1. The Member States of the European Union should continue to use the definitions (above and Table 1) established previously by European Commission (2009), incorporating any future amendments in these should they be agreed by the EU.
2. Other non-EU European countries, which typically are relatively ‘data-rich’ with respect to ornithological information, should adopt the EU definitions following review to ensure their applicability in the countries concerned.
3. Non-European countries (Africa, Middle East and western Eurasia) should establish definitions appropriate to the country concerned based on knowledge of species’ breeding biology in the appropriate regions. These should be used to determine the timing of the start and end of the breeding period. The following guidance may assist this process:
 - a. Review published knowledge of individual species breeding biology. Particularly valuable sources of information are given in Table 2.
 - b. Review any published knowledge, if this exists, of breeding biology in the country concerned.
 - c. Especially in the absence of published knowledge, consider what is known about the timing of breeding in neighbouring countries, or within the region, bearing in mind their position (north or south) of the country concerned and the implication that may have with respect to climatic/seasonal timing.

General recommendations related to intra-African migrants

As noted previously, intra-African migrants provide a range of problems related to the implementation of section 2 (and other parts) of AEWA’s Action Plan. The Technical Committee highlights the recommendations previously made by Dodman & Diagana (2006). These remain highly relevant to advancing conservation of such species:

1. Improve knowledge of the status of African waterbirds and their migratory patterns through:
 - Applied research of weather patterns, site conditions and waterbird seasonality;
 - Extending the African Waterbird Census (AfWC) to other seasons and other areas;

- Use/analysis of existing AfWC and other data to identify site linkages and migratory patterns;
 - Increased adoption of satellite telemetry;
 - Initial conservation focus on a series of “high profile species”;
 - Monitoring, research and conservation of threatened species;
 - Development of AFRING (African bird ringing scheme).
2. Identify key sites and site networks for intra-African migrants, especially threatened species.
 3. Develop Species Action Plans for African waterbirds.
 4. Promote increased focus on intra-African migrants in the implementation of the AEWA.
 5. Adopt a precautionary principle; it is often necessary to implement conservation action before knowing the full picture.
 6. Enhance awareness of African waterbirds, especially their values and ecological roles.
 7. Highlight the plight and lack of knowledge of threatened African waterbirds.
 8. Mobilise resources for conservation and monitoring of intra-African migrants, especially through development and subsequent implementation of a Conservation Strategy for African Waterbirds.

References

- Bauer, K.M. & Glutz von Blotzheim, U.N.** 1963. *Handbuch der Vögel Mitteleuropas, Volume 2.* Frankfurt am Main.
- Brown, L.H., Urban, E.K. & Newman, K.** (eds.) 1982. *The Birds of Africa, Volume I.* Academic Press, London. 521 pp.
- Cramp, S.** ed. 1985. *Handbook of the Birds of Europe, the Middle East and North Africa: the Birds of the Western Palearctic, Volume 4.* Oxford, Oxford University Press. 960 pp.
- Cramp, S. & Simmons, K.E.L.** (eds). 1977. *Birds of the Western Palearctic, Volume 1.* Oxford, Oxford University Press. 722 pp.
- Cramp, S. & Simmons, K.E.L.** eds. 1980. *Handbook of the Birds of Europe, the Middle East and North Africa: the Birds of the Western Palearctic, Volume 2.* Oxford, Oxford University Press. 695 pp.
- Cramp, S. & Simmons, K.E.L.** eds. 1983. *Handbook of the Birds of Europe, the Middle East and North Africa: the Birds of the Western Palearctic, Volume 3.* Oxford, Oxford University Press. 913 pp.
- Delany, S., Scott, D.A., Dodman, T., & Stroud, D.A.** (eds.) 2009. *An atlas of wader populations in Africa and western Eurasia.* Wetlands International, Wageningen, The Netherlands. 524 pp.
- del Hoyo, J., Elliott, A. & Sargatel, J.** (eds.) 1992. *Handbook of the Birds of the World, Volume 1. Hoatzin to Auks.* Lynx Edicions, Barcelona. 626 pp.
- del Hoyo, J., Elliott, A. & Sargatel, J.** (eds.) 1996. *Handbook of the Birds of the World, Volume 3. Ostrich to Ducks.* Lynx Edicions, Barcelona. 821 pp.
- Dodman, T. & Diagona, C.H.** 2006. Conservation dilemmas for intra-African migratory waterbirds. 218-223. In: *Waterbirds around the world.* Eds. G.C. Boere, C.A. Galbraith & D.A. Stroud. The Stationery Office, Edinburgh, UK. At: http://www.jncc.gov.uk/PDF/pub07_waterbirds_part3.4.1.pdf
- European Commission** 2009. *Key Concepts of Article 7(4) of Directive 79/409/EEC. Period of reproduction and pre-nuptial migration of Annex II bird species in the 27 EU Member States.* 476 pp. At: http://ec.europa.eu/environment/nature/conservation/wildbirds/hunting/key_concepts_en.htm
- Hockey, P.A.R., Dean, W.R.J. & Ryan, P.G.** (eds) 2005. *Roberts – Birds of Southern Africa. VIIth edition.* Cape Town. At: <http://www.robertsonline.co.za/portal.asp>
- Kear, J.** (ed.) 2005. *Ducks, geese and swans. Bird Families of the World XVI.* Two vols. Oxford University Press.
- Kushlan, J.A. & Hancock, J.A.** 2005. *The Herons. Bird Families of the World XVII.* Oxford University Press. 433 pp.
- Scott, D.A. & Rose, D.A.** 1996. *Atlas of Anatidae populations in Africa and western Eurasia.* Wetlands International Publication No. 41. Wageningen, The Netherlands.
- Taylor, B.** 1998. *Rails. A Guide to the Rails, Crakes, Gallinules and Coots of the World.* Christopher Helm/A & C Black.
- Urban, E.K., Fry, C.H. & Keith, S.** (eds.) 1986. *The Birds of Africa, Volume II.* Academic Press, London. 552 pp.
- Viljoen, P.J.** 2005. *AGRED's Gamebirds of South Africa: Field Identification and Management.* AGRED.

Table 1. Definitions for the start end of breeding periods as defined by the EU (European Commission 2009). Only species listed by AEWA are listed here.

Species	Start of breeding period	End of breeding period
<i>ANATIDAE</i>		
<i>Cygnus olor</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Anser fabalis</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Anser anser</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Anas penelope</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Anas strepera</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Anas crecca</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Anas platyrhynchos</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Anas acuta</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Anas querquedula</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Anas clypeata</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Netta rufina</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Aythya ferina</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Aythya fuligula</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Aythya marila</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Somateria mollissima</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds

Species	Start of breeding period	End of breeding period
<i>Clangula hyemalis</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Melanitta nigra</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Melanitta fusca</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Bucephala clangula</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Mergus serrator</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Mergus merganser</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
RALLIDAE		
<i>Rallus aquaticus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds (3 decades ⁴ after hatching)
<i>Gallinula chloropus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds (5 decades after hatching)
<i>Fulica atra</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds (6 decades after hatching)
HAEMATOPODIDAE		
<i>Haematopus ostralegus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
CHARADRIIDAE		
<i>Pluvialis apricaria</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Vanellus vanellus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
SCOLOPACIDAE		
<i>Philomachus pugnax</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Lymnocyptes minimus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds (c. 4 decades after hatching)
<i>Gallinago gallinago</i>	<ul style="list-style-type: none"> • occupation of the breeding sites with courtship display • construction of the nest in all other cases 	full flight of young birds (c. 4 decades after hatching)

⁴ i.e. periods of ten days

Species	Start of breeding period	End of breeding period
<i>Scolopax rusticola</i>	<ul style="list-style-type: none"> • occupation of the breeding sites (roding) 	full flight of young birds (c. 4 decades after hatching)
<i>Limosa limosa</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Limosa lapponica</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Numenius phaeopus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Numenius arquata</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Tringa erythropus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
<i>Tringa totanus</i>	<ul style="list-style-type: none"> • occupation of the breeding sites where it is mainly migratory • construction of the nest in all other cases 	full flight of young birds
<i>Tringa nebularia</i>	<ul style="list-style-type: none"> • occupation of the breeding sites 	full flight of young birds
LARIDAE		
<i>Larus ridibundus</i>	<ul style="list-style-type: none"> • courtship display at breeding sites (2 decades before egg laying) 	full flight of young birds
<i>Larus canus</i>	<ul style="list-style-type: none"> • courtship display at breeding sites (2 decades before egg laying) 	full flight of young birds
<i>Larus fuscus</i>	<ul style="list-style-type: none"> • courtship display at breeding sites (3 decades before egg laying) 	full flight of young birds
<i>Larus argentatus</i>	<ul style="list-style-type: none"> • courtship display at breeding sites (3 decades before egg laying) 	full flight of young birds
<i>Larus cachinnans</i>	<ul style="list-style-type: none"> • courtship display at breeding sites (3 decades before egg laying) 	full flight of young birds
<i>Larus marinus</i>	<ul style="list-style-type: none"> • courtship display at breeding sites (3 decades before egg laying) 	full flight of young birds

Table 2. Valuable sources of information useful for determining the start and end of breeding seasons and the timing of migration, of African and Eurasian waterbirds.

Waterbird family	Information on breeding seasons	Information on migration periods
<i>Sphenisciformes</i> <i>Spheniscidae</i> Penguins	Africa: Brown <i>et al.</i> (1982); Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)	Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)
<i>Gaviiformes</i> <i>Gaviidae</i> Divers	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)
<i>Podicipediformes</i> <i>Podicipedidae</i> Grebes	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)
<i>Pelacaniformes</i> <i>Phaethontidae</i> Tropicbirds <i>Pelecanidae</i> Pelicans <i>Sulidae</i> Gannets and boobies <i>Phalacrocoracidae</i> Cormorants <i>Fregatidae</i> Frigatebirds	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)
<i>Ciconiiformes</i> <i>Ardeidae</i> Herons and Egrets <i>Ciconiidae</i> Storks <i>Balaenicipitidae</i> Shoebill <i>Threskiornithidae</i> Ibises	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992); Kushlan & Hancock (2005)	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992); Kushlan & Hancock (2005)
<i>Phoenicopteriformes</i> <i>Phoenicopteridae</i> Flamingos	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1992)
<i>Anseriformes</i> <i>Anatidae</i> Ducks, geese and swans	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005); Viljoen (2005)	Western Palearctic: Cramp & Simons (1977) Africa: Brown <i>et al.</i> (1982) ; Hockey <i>et al.</i> (2005); Viljoen (2005)

Waterbird family	Information on breeding seasons	Information on migration periods
	All regions: Bauer & Glutz von Blotzheim (1963); del Hoyo <i>et al.</i> (1992); Scott & Rose (1996); Kear (2005)	All regions: del Hoyo <i>et al.</i> (1992); Scott & Rose (1996); Kear (2005)
<i>Gruiformes</i> <i>Gruidae</i> Cranes <i>Rallidae</i> Rails, flufftails, crakes, gallinules, moorhens and coots	Western Palearctic: Cramp & Simons (1980) Africa: Urban <i>et al.</i> (1986) ; Hockey <i>et al.</i> (2005)); Viljoen (2005) All regions: del Hoyo <i>et al.</i> (1996) Rallidae: Taylor (1998)	Western Palearctic: Cramp & Simons (1980) Africa: Urban <i>et al.</i> (1986) ; Hockey <i>et al.</i> (2005)); Viljoen (2005) All regions: del Hoyo <i>et al.</i> (1996) Rallidae: Taylor (1998)
<i>Charadriiformes</i> <i>Dromadidae</i> Crab Plover <i>Haematopodidae</i> Oystercatchers <i>Recurvirostridae</i> Stilts and avocets <i>Burhinidae</i> Thick-knees <i>Glareolidae</i> Practincoles <i>Charadriidae</i> Plovers <i>Scolopacidae</i> Snipes, curlews and sandpipers <i>Stercorariidae</i> Skuas <i>Laridae</i> Gulls <i>Sternidae</i> Terns <i>Rynchopidae</i> Skimmers <i>Alcidae</i> Auks	Western Palearctic: Cramp & Simmons (1983); Cramp (1985) Africa: Urban <i>et al.</i> (1986) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1996)	Western Palearctic: Cramp & Simmons (1983); Cramp (1985) Africa: Urban <i>et al.</i> (1986) ; Hockey <i>et al.</i> (2005) All regions: del Hoyo <i>et al.</i> (1996) All regions for waders: Delany <i>et al.</i> (2009)