

*Secretariat provided by the
United Nations Environment
Programme (UNEP)*

*Doc: AEWA/Inf. 3.1
Date: 14 October 2005
Original: English*

AGREEMENT ON THE CONSERVATION OF
AFRICAN-EURASIAN MIGRATORY WATERBIRDS

**Report on the Conservation Status of Migratory
Waterbirds in the Agreement Area¹**

Third Edition

Draft Report prepared by Wetlands International

Simon Delany

September 2005

¹ This is a preliminary version of the report containing only tables. Full version will be provided before MOP3.

TABLE 1: POPULATION SIZES AND TRENDS OF WATERBIRD SPECIES INCLUDED IN THE AGREEMENT

KEY TO COLUMN HEADINGS:

Population

The migratory populations of the waterbird species covered under this Agreement. These populations may be the entire population of a species, the entire population of a distinct subspecies, a discrete biogeographical population of a monotypic species or of a subspecies, or that 'population' of birds which breeds and/or spends the boreal winter in one or more of the major sub-regions of the African-Eurasian region. For an explanation of the ways in which populations are listed, see section 2.1 in the second (2002) edition of this report.

Population size

Population estimates given in the column headed 'CSR 1999' are taken from the first edition of the *Report on the Conservation Status of Migratory Waterbirds in the Agreement Area* (Wetlands International 2000). Those in the second column, headed 'CSR 2002' are taken from the second edition of the *Report on the Conservation Status of Migratory Waterbirds in the Agreement Area* (Wetlands International 2002).

In all cases, the numerical estimates are given as number of individuals, although in many cases the estimates have been derived from numbers of breeding pairs. In these latter estimates, the total number of individuals has been obtained by multiplying the number of breeding pairs by a factor of three. For many species and populations, only a rough indication of population size is available. In these cases, the size of the population is given in one of five categories:

- A: Less than 10,000 individuals
- B: 10,000-25,000 individuals
- C: 25,000-100,000 individuals
- D: 100,000-1,000,000 individuals
- E: Over 1,000,000 individuals

Trends

Information on trends is confined to the situation in recent decades. Brackets are used to indicate either that there is some doubt about the trend, or that a particular trend has been reported in only a part of the species' range, and may not be applicable throughout most of its range. The trends given in the column headed 'Trend 1999' are taken from the first edition of the *Report on the Conservation Status of Migratory Waterbirds in the Agreement Area*. Those in the second column, headed 'Trend 2002' are taken from the second edition of the *Report on the Conservation Status of Migratory Waterbirds in the Agreement Area* (Wetlands International 2002).

INC	-	Increasing	DEC	-	Decreasing
STA	-	Stable	?	-	Unknown

Sources

Sources of information included since the 2002 edition of this report are numbered in the table, and listed at the end.

All information which has not appeared in earlier editions of this table is shaded in this edition.

Note: The Tables in the AEWA Conservation Status Report were updated with information compiled during 2005.

Table 1, headed “**Population sizes and trends of waterbird species included in the Agreement**” includes details of the status of waterbirds compiled for AEWA MOP 1 in 1999 and AEWA MOP 2 in 2002. The proposed revisions of population estimates and trends compiled in 2005 are included in two new columns, which are shaded grey. Information is included in the new columns even if the estimate or trend have not changed, so that these new columns provide all the current information on population status. Sources of changes in the population estimates and trends are given in the form of a code number in two additional new shaded columns (one for estimates, the other for trends). The 100 new sources are all listed at the end of Table 1, together with the appropriate code number, in numerical and alphabetical order. The easiest way to identify changes in the table is to look at the rows where there is an entry in the source column. Changes in taxonomic treatment and nomenclature are also highlighted with grey shading. A few single populations have been split into two. These have their names shaded in grey, and only include information on populations, trends and sources in the new, shaded columns. The previous treatment, when these populations were combined in the 1999 and 2002 reviews, appears on the line above the updated treatment, and information in these cases only appears in the old, unshaded columns. One comment about the substance of this table might save confusion: it is in the nature of these population estimates that estimated numbers sometimes increase as knowledge of populations improves. For a number of populations, the estimate has increased during the assessments in 2002 and 2005, even though it is known that the population may be stable or decreasing.

Table 2, headed “**Population sizes and trends of waterbird species proposed for inclusion in the Agreement**” includes 33 populations of 21 species being considered for inclusion in the Agreement. The data in this table were compiled by the Technical Committee, and updated for three species (*Sterna anaethetus*, *Anous minutus* and *Anous tenuirostris*) in September 2005.

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
SPHENISCIDAE								
<i>Spheniscus demersus</i>								
- <i>Southern Africa</i>	-	180,000	180,000			DEC	DEC	
GAVIIDAE								
<i>Gavia stellata</i>								
- Northwestern Europe (win)	D	183,000-420,000	150,000-450,000	7	DEC	DEC	DEC	7
- Caspian, Black Sea & E Med (win)	?	?	?		?	?	?	
<i>Gavia arctica arctica</i>								
- N Europe & Western Siberia/Europe	120,000	360,000-690,000	250,000-500,000	7	STA	DEC	DEC	7
<i>Gavia arctica suschkini</i>								
- Central Siberia/Caspian	?	?	?		?	?	?	
<i>Gavia immer</i>								
- Europe (win)	5,000	5,000	5,000		?	?	?	
<i>Gavia adamsii</i>								
- Northern Europe (win)	A/B	A	A		?	?	?	
PODICIPEDIDAE								
<i>Tachybaptus ruficollis ruficollis</i>								
- Europe & Northwest Africa	D	230,000-450,000	300,000-510,000	7	STA	STA	STA	7
<i>Podiceps cristatus cristatus</i>								
- Northwest & Western Europe	D	368,000-579,000	290,000-420,000	7		INC	DEC	7
- Black Sea & Mediterranean (win)	D	>600,000	580,000-870,000	7		INC	DEC	7
- Caspian & Southwest Asia (win)	10,000	10,000	10,000			?	?	
<i>Podiceps cristatus infuscatus</i>								
- Eastern Africa (Ethiopia to N Zambia)	<1,000	<1,000	<1,000			DEC	DEC	
- Southern Africa	A	A	A			INC	INC	
<i>Podiceps grisegena grisegena</i>								
- Northwestern Europe (win)	C	C	42,000-60,000	7	STA	STA	DEC	7
- Black Sea & Mediterranean (win)	C	C	41,000-107,000	7	?	STA	DEC	7
- Caspian (win)	15,000	15,000	15,000		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Podiceps auritus auritus</i>								
- NW Europe (large billed)	5,000	2,600-4,100	4,600-6,800	7	STA/INC	STA/INC	STA	7
- NE Europe (small billed)	C	C	14,200-26,000	7	(DEC)	(STA)	DEC	7
- Caspian & South Asia (win)	B	B	B		?	?	?	
<i>Podiceps nigricollis nigricollis</i>								
- Europe/S & W Europe & N Africa	100,000	117,000-450,000	159,000-268,000	7	(STA)	STA/INC	DEC	7
- Western Asia/SW & South Asia	25,000	25,000	25,000		INC	INC	INC	
<i>Podiceps nigricollis gurneyi</i>								
- Southern Africa	B/C	10,000-20,000	10,000-20,000		INC	INC	INC	
PELECANIDAE								
<i>Pelecanus onocrotalus</i>								
- Southern Africa	-	18,000	30,000	81 22	-	(STA)	INC	81
- West Africa	30,000	60,000	60,000		?	STA	INC	71
- Eastern Africa	-	150,000	140,000	14 22	-	STA	STA	
- Europe & Western Asia (bre)	70,000	20,000-33,000	20,000-33,000		DEC	DEC	DEC	
<i>Pelecanus rufescens</i>								
- Tropical Africa & SW Arabia	D	50,000-100,000	50,000-100,000		STA	STA	STA	
<i>Pelecanus crispus</i>								
- Black Sea & Mediterranean (win)	2,000-3,000	2,300-3,200	4,350-4,800	15 75	STA	STA	INC	75
- SW Asia & S Asia (win)	10,000-13,000	10,000-12,500	6,000-9,000	15 75	STA	STA	DEC	75
SULIDAE								
<i>Sula (Morus) capensis</i>								
- Southern Africa	-	346,000	346,000		-	DEC	DEC	
PHALACROCORACIDAE								
<i>Phalacrocorax coronatus</i>								
- Coastal Southwest Africa	5,330	8,700	8,700		?	STA	STA	
<i>Phalacrocorax pygmeus</i>								
- Black Sea & Mediterranean	25,000	23,000-37,000	23,000-37,000		STA/INC	STA/INC	STA/INC	
- Southwestern Asia	C	C	C		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Phalacrocorax neglectus</i>								
- Coastal Southwest Africa	18,000	9,700	11,100	81 36	?	DEC	DEC	
<i>Phalacrocorax carbo carbo</i>								
- Northwest Europe	120,000	114,000	120,000	7	INC	INC	INC	7
<i>Phalacrocorax carbo sinensis</i>								
- Northern & Central Europe	200,000	275,000-340,000	380,000-405,000	7	INC	INC	INC	7
- Black Sea & Mediterranean	100,000	>130,000-160,000	350,000-450,000	7	INC	INC	INC	7
- West & Southwest Asia	100,000	100,000	100,000		?	?	?	
<i>Phalacrocorax carbo lucidus</i>								
Coastal West Africa			35,000	21			STA	21
Central & Eastern Africa	D	135,000-535,000	200,000-500,000	21	?	STA	STA	
- Southern Africa	B	11,000-13,000	13,000	36	?	STA	STA	
<i>Phalacrocorax nigrogularis</i>								
Persian Gulf & Arabian Sea	500,000-1,000,000	450,000-750,000			(DEC)	DEC		
Arabian Coast			270,000	44			DEC	9 44
Gulf of Aden, Socotra, Arabian Sea			60,000	43 9			STA/IN C	22
<i>Phalacrocorax capensis</i>								
	550,000	216,000	300,000	9 57 22	?	DEC	DEC	
ARDEIDAE								
<i>Egretta ardesiaca</i>								
- Subsaharan Africa	B/C	B/C	C	38 85	DEC	DEC	STA	22
<i>Egretta vinaceigula</i>								
- South-central Africa	5,000-10,000	3,000-5,000	3,000-5,000		DEC	DEC	DEC	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Egretta garzetta garzetta</i>								
- Subsaharan Africa	?	100,000-500,000	200,000-500,000	21	?	?		
- Europe, Black Sea & Med/W & C Africa	100,000-150,000	140,000-210,000			INC	INC		
W Europe, NW Africa			125,000-143,000	32			INC	
C&E Europe, Black Sea, E Mediterranean			44,000-72,400	32			STA	32
- Western Asia/SW Asia, NE & E Africa	C	C	C		?	?	?	
<i>Egretta gularis gularis</i> *								
- West Africa	?	B/C	B/C		?	(STA)	STA	21
<i>Egretta gularis schistacea</i> *								
- NE Africa & Red Sea	?	B/C	B/C		?	?	?	
- SW Asia & South Asia	17,000	17,000	17,000		?	?	?	
<i>Egretta dimorpha</i> *								
- Coastal Eastern Africa	?	10,000	10,000		?	STA	STA	
<i>Ardea cinerea cinerea</i>								
- Subsaharan Africa	?	D	D		?	STA	STA	
- Europe & North Africa (bre)	400,000-500,000	480,000-600,000			INC	INC		
N & W Europe			263,000-286,000	54			INC	
C & E Europe			189,000-256,000	54 32			INC	
- West & SW Asia (bre)	B/C	C/D	C/D		?	STA/DEC	?	
<i>Ardea melanocephala</i>								
- Subsaharan Africa	D/E	100,000-500,000	100,000-500,000		INC	INC	INC	
<i>Ardea purpurea purpurea</i>								
- Tropical Africa	C	75,000-100,000	75,000-100,000		?	STA	STA	
- W Europe & W Mediterranean/W Africa	B	12,000-13,200	11,500-12,100	54	DEC	DEC	DEC	75
- E Europe & SW Asia/Subsaharan Africa	D	D	135,000-300,000	75	(DEC)	DEC	DEC	7
<i>Casmerodius albus albus</i>								
- W, C & SE Europe/Black Sea & Med	7,000-17,000	12,000-22,500	38,800-54,300	75	STA/INC	INC	INC	75
- West Asia/Southwest Asia	B/C	C	C		?	(INC)	(INC)	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Casmerodius albus melanorhynchos</i>								
- Subsaharan Africa & Madagascar	C	100,000-500,000	100,000-500,000		?	STA	STA	
<i>Mesophoyx intermedia brachyrhyncha</i>								
- Subsaharan Africa	?	C	C		?	STA	STA	
<i>Bubulcus ibis ibis</i>								
- Southern Africa	D	D	D		?	INC	INC	
- Tropical Africa	E	E	E		?	?	?	
- SW Europe & NW Africa	>250,000	300,000-450,000			INC	INC		
SW Europe			250,000-310,000	75			INC	75
NW Africa			100,000-150,000	32			INC	79
- East Mediterranean & SW Asia	A/B	A/B	B/C	75	?	?	INC	75
<i>Ardeola ralloides ralloides</i>								
- Med., Black S. & N Africa/Subs. Africa	-	40,000-70,000			-	DEC		
SW Europe, NW Africa (bre)			2,700-5,600	75			INC	75
C & E Europe/Black Sea & E Mediterranean (bre)			42,000-76,000	75			DEC	75
- West & SW Asia/Subsaharan Africa ²	-	C	C		-	?	?	
<i>Ardeola ralloides paludivaga</i>								
- Subsaharan Africa & Madagascar	C	100,000-500,000	300,000-600,000	22 85	?	STA	STA	
<i>Ardeola idae</i>								
- Madagascar & Aldabra/Cen & E Africa	5,000	2,000-6,000	2,000-6,000		DEC	DEC	DEC	
<i>Ardeola rufiventris</i>								
- Tropical Eastern & Southern Africa	B/C	B/C	B/C		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Nycticorax nycticorax nycticorax</i>								
- Subsaharan Africa & Madagascar	C	C/D	C/D		?	STA	STA	
- Europe & NW Africa/Med & Africa	100,000-200,000	150,000-230,000			DEC	DEC		
W Europe, NW Africa (bre)			61,000-97,000	54			DEC	
C & E Europe, Black Sea, E Mediterranean (bre)			92,100-138,000	75			DEC	
- Western Asia/SW Asia & NE Africa	C	C	C		?	?	?	
<i>Ixobrychus minutus minutus</i>								
- Europe & N Africa/Subsaharan Africa	D	D			DEC	DEC		
W Europe, NW Africa/Subsaharan Africa			11,900-17,900	54			DEC	75
C & E Europe, Black Sea & E Mediterranean/Subsaharan Africa			110,000-325,000	75			DEC	75
- West & SW Asia/Subsaharan Africa	C	C	C		?	?	?	
<i>Ixobrychus minutus payesii</i>								
- Subsaharan Africa	?	C	C		?	?	?	
<i>Ixobrychus sturmii</i>								
- Subsaharan Africa	B/C	C	C		?	?	?	
<i>Botaurus stellaris stellaris</i>								
- Europe (bre)	C	C			DEC	DEC		
W Europe, NW Africa (bre)			5,880-6,730	54			DEC	75
C & E Europe, Black Sea & E Mediterranean (bre)			53,800-124,200	54			DEC	75
- Southwestern Asia (win)	A/B	A/B	C	75	?	?	?	75
<i>Botaurus stellaris capensis</i>								
- Southern Africa	?	5,000	5,000		?	DEC	DEC	
CICONIIDAE								
<i>Mycteria ibis</i>								
- Subsaharan Africa (excl. Madagascar)	C	50,000-100,000	75,000-100,000	22	STA	STA	STA	10

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Anastomus lamelligerus lamelligerus</i>								
- Subsaharan Africa	C/D	400,000-800,000	300,000-500,000	21	STA	STA	STA	
<i>Ciconia nigra</i>								
- Southern Africa	<1,500	2,850-4,740	1,560-4,050	21	STA	STA	(STA)	21
- SW Europe/West Africa	1,000	1,300-1,370	1,300-1,370		STA/IN C	INC	INC	
- Central & E Europe/Subsaharan Africa	20,000-30,000	19,500-28,000	19,500-28,000		?	DEC	DEC	
<i>Ciconia abdimii</i>								
- Subsaharan Africa & SW Arabia	D	300,000-600,000	300,000-600,000		STA	(DEC)	DEC	21
<i>Ciconia episcopus microscelis</i>								
Sub-Saharan Africa	C	B/C	B/C		?	?	?	
<i>Ciconia ciconia ciconia</i>								
- Southern Africa	30	24	20	91	INC	STA	STA	91
W Europe & NW Africa/Subsaharan Africa	100,000	93,000	93,000		STA/IN C	INC	INC	
- Central & E Europe/Subsaharan Africa	400,000	390,000-400,000	390,000-400,000		DEC	STA	STA	
- West Asia/Southwest Asia	C	B	B		?	DEC	DEC	
<i>Leptoptilos crumeniferus</i>								
- Subsaharan Africa	100,000	100,000-300,000	200,000-500,000	22	INC	INC	INC	
BALAENICIPITIDAE								
<i>Balaeniceps rex</i>								
- Central Tropical Africa	12,000-15,000	5,000-10,000	5,000-8,000	21	?	DEC	DEC	
THRESKIORNITHIDAE								
<i>Plegadis falcinellus falcinellus</i>								
- Subsaharan Africa (bre)	?	1.0-2.0 million	1.0-2.0 million		?	?	?	
- Black Sea & Med/West Africa	40,000-50,000	49,000-57,000	48,000-66,000	7	DEC	DEC	DEC	7
- Southwestern Asia/E Africa	C	C	C		?	?	?	
<i>Geronticus eremita</i>								
- Morocco	200	190	227	8	DEC	STA	DEC	8
- Southwestern Asia	>27	>27	7	6	DEC	DEC	DEC	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Threskiornis aethiopicus aethiopicus</i>								
- Subsaharan Africa	D	200,000-450,000	200,000-450,000		STA	STA	STA	
- Iraq & Iran	200	200	200		DEC	DEC	DEC	
<i>Platalea leucorodia leucorodia</i>								
- West Europe/W Medit. & West Africa	6,500	9,945	11,300	61 62	INC	INC	INC	61
- C & SE Europe/Med & Tropical Africa	5,000-15,000	11,670	11,670		DEC	DEC	DEC	
<i>Platalea leucorodia archeri</i>								
- Red Sea & Somalia	500-1,500	1,250	1,500-2,250	39 22	?	?	DEC	39
<i>Platalea leucorodia balsaci</i>								
- Coastal West Africa (Mauritania)	5,000-6,000	7,000	6,000-7,000	61	?	STA	STA	
<i>Platalea leucorodia major</i>								
- W Asia/SW & S Asia	23,000	23,000	23,000		?	?	?	
<i>Platalea alba</i>								
- Subsaharan Africa	A/B	B/C	B/C		?	STA	STA	
PHOENICOPTERID AE								
<i>Phoenicopterus roseus</i>								
- West Africa	40,000	40,000	30,000	20 3 4	?	STA	FLU	20 3 4
- Eastern Africa	35,000	35,000	35,000		DEC	STA	STA	
- Southern Africa (to Madagascar)	55,000	65,000-87,000	65,000-87,000		DEC	STA	STA	
- Western Mediterranean	80,000	100,000	165,000	3	INC	INC	INC	3
East Mediterranean, SW & S Asia	500,000	290,000			STA	STA		
East Mediterranean			60,000	4			?	
SW & S Asia			325,000	63 28			?	
<i>Phoenicopterus minor</i>								
- West Africa	15,000	15,000	15,000-25,000	87 90 23	STA	STA	STA	87 90 23
- Eastern Africa	4,000,000	2.0-4.0 million	1.5-2.5 million	75	DEC	DEC	DEC	
- Southern Africa (to Madagascar)	40,000	55,000-65,000	55,000-65,000		DEC	STA	STA	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
ANATIDAE								
<i>Dendrocygna bicolor</i>								
- West Africa (Senegal to Chad)	100,000	100,000	50,000-100,000	65 85	?	?	?	
- Eastern & Southern Africa	200,000-500,000	150,000-350,000	150,000-350,000	21	?	?	?	
<i>Dendrocygna viduata</i>								
- West Africa (Senegal to Chad)	250,000	250,000-500,000	600,000-700,000	88 22	INC	INC	INC	
- Eastern & Southern Africa	1.0-2.0 million	D	D		INC	INC	INC	
<i>Thalassornis leuconotus leuconotus</i>								
- West Africa	1,000	<1,000	<500	22	DEC	DEC	DEC	
- Eastern & Southern Africa	10,000-25,000	10,000-25,000	10,000-25,000		STA	STA	STA	
<i>Oxyura leucocephala</i>								
- West Mediterranean (Spain & Morocco)	1,200	2,000-4,500	2,500	82	INC	INC	INC	
- Algeria & Tunisia	400	400	400-600	1	STA	STA	STA	
- E Mediterranean, Turkey & SW Asia	8,000-15,000	8,000-15,000	5,000-10,000	50	DEC	DEC	DEC	
<i>Oxyura maccoa</i>								
- Eastern Africa	15,000-25,000	1,500	1,000-1,500	21	DEC	DEC	DEC	
- Southern Africa	15,000-25,000	A	A		INC	INC	INC	
<i>Cygnus olor</i>								
- NW Mainland & Central Europe	210,000	210,000	250,000	96	INC	INC	INC	
- Black Sea	45,000	45,000	45,000		INC	INC	INC	
- W & Central Asia/Caspian	250,000	250,000	250,000		INC	INC	INC	
<i>Cygnus cygnus</i>								
- Iceland/UK & Ireland	16,000	20,900	20,900		?	INC	INC	
- NW Mainland Europe	59,000	59,000	59,000		INC	INC	INC	
- N Europe & W Sib/B Sea & E Med	17,000	17,000	17,000		DEC	DEC	DEC	
- W & Cent Siberia/Caspian	20,000	20,000	20,000		DEC	DEC	DEC	
<i>Cygnus columbianus bewickii</i>								
- W Siberia & NE Europe/NW Europe	29,000	29,000	20,000	68 69	INC	DEC	DEC	68 69
- Northern Siberia/Caspian	500	500	1,000	77	?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Anser brachyrhynchus</i>								
- E Greenland & Iceland/UK	250,000	241,000	270,000	99	INC	INC	INC	
- Svalbard/NW Europe	37,000	37,000	42,000	53	INC	STA	INC	53
<i>Anser fabalis fabalis</i>								
- NE Europe/NW Europe	100,000	100,000	70,000-90,000	60	STA	STA	DEC	60
<i>Anser fabalis rossicus</i>								
- W & C Siberia/NE & SW Europe	600,000	600,000	600,000		?	?	?	
<i>Anser fabalis johanseni</i>								
- W & C Sib/Turkmenistan to W China	-	?	?		-	?	?	
<i>Anser albifrons albifrons</i>								
- NW Siberia & NE Europe/NW Europe	600,000	600,000	1,000,000	28	INC	(INC)	STA	47
- Western Siberia/Central Europe	100,000	100,000	10,000-40,000	52	DEC	DEC	DEC	
- Western Siberia/Black Sea & Turkey	650,000	650,000	350,000-700,000	52	(STA)	STA	STA	
- Northern Siberia/Caspian & Iraq	15,000	15,000	15,000		DEC	DEC	DEC	
<i>Anser albifrons flavirostris</i>								
- Greenland/Ireland & UK	33,000	30,000	27,000	26	INC	STA	DEC	26
<i>Anser erythropus</i>								
- N Europe & W Sib/B Sea & Caspian	15,000	8,000-13,000	8,000-13,000		DEC	DEC	DEC	
<i>Anser anser anser</i>								
- Iceland/UK & Ireland	80,000	89,100	87,200	99	DEC	(STA)	(STA)	
- NW Europe/SW Europe	200,000	400,000	500,000	75	INC	INC	INC	
- Central Europe/N Africa	25,000	25,000	25,000		INC	INC	INC	
<i>Anser anser rubrirostris</i>								
- Black Sea & Turkey	85,000	85,000	85,000		?	?	?	
- Western Siberia/Caspian & Iraq	>100,000	>100,000	250,000	73	INC	INC	INC	
<i>Branta leucopsis</i>								
- E Greenland/Scotland & Ireland	40,000	54,100	56,400	97	INC	INC	INC	
- Svalbard/SW Scotland	23,000	23,000	27,000	98	INC	INC	INC	
- Russia/Germany & Netherlands	267,000	360,000	420,000	75	INC	INC	INC	
<i>Branta bernicla bernicla</i>								
- Western Siberia/Western Europe	300,000	190,000	200,000	24	INC/ST A	DEC	DEC	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Branta bernicla hrota</i>								
- Svalbard/Denmark & UK	5,000	5,000	7,000	17 11	STA	STA	INC	
- Canada & Greenland/Ireland	20,000	20,000	26,400	41	STA	STA	STA	
<i>Branta ruficollis</i>								
- Northern Siberia/Black Sea & Caspian	70,000	88,000	38,500	18	STA/IN C	INC	DEC	18
<i>Alopochen aegyptiacus</i>								
- West Africa	10,000-25,000	10,000-25,000	5,000-10,000	85 22	?	DEC	DEC	
- Eastern & Southern Africa	200,000-500,000	200,000-500,000	200,000-500,000		?	STA	STA	
<i>Tadorna ferruginea</i>								
- Northwest Africa	3,000	3,000	3,000		DEC	DEC	DEC	
- E Mediterranean & Black Sea/NE Africa	20,000	20,000	20,000		DEC	DEC	DEC	
- Western Asia & Caspian/Iran & Iraq	35,000	35,000	50,000	28	INC	INC	INC	
<i>Tadorna cana</i>								
- Southern Africa	42,000	42,000	50,000	33	STA	STA	INC	35
<i>Tadorna tadorna</i>								
- Northwestern Europe	300,000	300,000	300,000		INC	STA	STA	
- Black Sea & Mediterranean	75,000	75,000	75,000		STA/DEC	DEC	DEC	
- Western Asia/Caspian & Middle East	80,000	80,000	80,000		INC	INC	INC	
<i>Plectropterus gambensis gambensis</i>								
- West Africa	50,000	100,000	50,000-100,000	22 85	DEC	STA	STA	
- Eastern Africa (Sudan to Zambia)	200,000-300,000	200,000-300,000	200,000-300,000		STA	STA	STA	
<i>Plectropterus gambensis niger</i>								
- Southern Africa	50,000-100,000	50,000-100,000	50,000-100,000		STA	INC	INC	
<i>Sarkidiornis melanotos melanotos</i>								
- West Africa	50,000	50,000-100,000	50,000-80,000		STA	STA	DEC	85
- Southern & Eastern Africa	500,000-1,000,000	100,000-500,000	100,000-500,000		STA	STA	STA	
<i>Nettapus auritus</i>								
- West Africa	20,000-30,000	<10,000	<10,000		?	DEC	DEC	
- Southern & Eastern Africa	100,000-250,000	100,000-250,000	100,000-250,000		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Anas capensis</i>								
Eastern Africa (Rift Valley)	A/B	5,000-15,000	5,750-7,000	2	STA	DEC	STA	2
Lake Chad Basin			<500	2			DEC	
- Southern Africa (N to Angola & Zambia)	100,000-250,000	100,000-250,000	100,000-250,000		INC	INC	INC	
<i>Anas strepera strepera</i>								
- Northwestern Europe	30,000	50,000	60,000	28	INC	INC	INC	
- NE Europe/Black Sea & Med	75,000-150,000	75,000-150,000	75,000-150,000		DEC	DEC	(STA)	96
- W Siberia/SW Asia & NE Africa	130,000	130,000	130,000		?	?	?	
<i>Anas penelope</i>								
- W Siberia & NE Europe/NW Europe	1,250,000	2,000,000	1,500,000	96	INC	INC	STA	96
- W Siberia & NE Europe/B Sea & Med	560,000	300,000	300,000		DEC	DEC	DEC	
- Western Siberia/SW Asia & NE Africa	250,000	250,000	250,000		DEC	DEC	DEC	
<i>Anas platyrhynchos platyrhynchos</i>								
- Northwestern Europe	5,000,000	4,500,000	4,500,000		STA	DEC	DEC/STA	
- N Europe/Western Mediterranean	1,000,000	1,000,000	1,000,000		INC	(STA)	(STA)	
- E Europe/Black Sea & E Mediterranean	2,250,000	2,000,000	2,000,000		DEC	DEC	(STA)	
- Western Siberia/SW Asia	800,000	800,000	800,000		?	?	?	
<i>Anas undulata undulata</i>								
- Southern Africa	>100,000	>100,000	>100,000		STA	STA	STA	
<i>Anas clypeata</i>								
- NW & Central Europe (win)	40,000	40,000	40,000		STA	(STA)	STA	96
- W Sib, NE & E Eur/S Eur & W Africa	450,000	450,000	450,000		(DEC)	?	STA	96
- W Siberia/SW Asia, NE & E Africa	400,000	400,000	400,000		DEC	(DEC)	(DEC)	
<i>Anas erythrorhyncha</i>								
- Southern Africa	500,000-1,000,000	500,000-1,000,000	500,000-1,000,000		STA	STA	STA	96
- Eastern Africa	100,000-300,000	100,000-300,000	100,000-300,000		STA	STA	STA	
- Madagascar	15,000-25,000	15,000-25,000	15,000-25,000		DEC	DEC	DEC	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Anas acuta</i>								
- Northwestern Europe	60,000	60,000	60,000		DEC	DEC	STA	96
- W Sib, NE & E Eur/S Eur & W Africa	1,200,000	1,000,000	750,000	85 16	DEC	DEC	DEC	
- W Siberia/SW Asia & Eastern Africa	700,000	700,000	700,000		?	?	?	
<i>Anas querquedula</i>								
- W Siberia & Europe/West Africa	2,000,000	2.0-3.3 million	2.0 million	85 7	DEC	DEC	DEC	
- W Siberia/SW Asia, NE & East Africa	100,000-200,000	100,000-200,000	100,000-200,000		?	?	?	
<i>Anas crecca crecca</i>								
- Northwestern Europe	400,000	400,000	500,000	96	INC	STA	INC	96
- W Siberia & NE Europe/B Sea & Med	750,000-1,375,000	1,000,000	750,000-1,375,000		STA	(STA)	(STA)	
- W Siberia/SW Asia & NE Africa	1,500,000	1,500,000	1,500,000		DEC	DEC	DEC	
<i>Anas hottentota</i>								
- Lake Chad Basin	5,000-10,000	<1,000	1,000-5,000	21	DEC	DEC	DEC	
- Eastern Africa (south to N Zambia)	100,000-300,000	C	C		STA	STA	STA	
- Southern Africa (north to S Zambia)	C	C	C		STA	STA	STA	
<i>Marmaronetta angustirostris</i>								
- W Mediterranean/W Med & West Africa	3,000	3,000-5,000	3,000-5,000		DEC	?	FLU	30
- Eastern Mediterranean	1,000	1,000	1,000		DEC	DEC	DEC	
- Southwestern Asia	5,000-15,000	5,000-15,000	5,000-15,000		DEC	DEC	DEC	
<i>Netta rufina</i>								
- SW & Cent Europe/W Mediterranean	25,000	50,000	50,000		INC	INC	INC	
- Black Sea & E Mediterranean	50,000	20,000-43,500	20,000-43,500		DEC	DEC	DEC	
- Western & Central Asia/SW Asia	200,000	250,000	250,000		STA	(STA)	(STA)	
<i>Netta erythrophthalma brunnea</i>								
- Southern & Eastern Africa	30,000-70,000	30,000-70,000	30,000-70,000		STA	STA	STA	
<i>Aythya ferina</i>								
- NE Europe/NW Europe	350,000	350,000	350,000		DEC	STA	DEC	96
- Central & NE Europe/B Sea & Medit.	1,000,000	1,100,000	1,000,000	96	DEC	INC	DEC	96
- Western Siberia/SW Asia	350,000	350,000	350,000		(DEC)	(DEC)	(DEC)	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Aythya nyroca</i>								
- W Mediterranean/North & West Africa	2,000-3,000	2,000-3,000	2,400-2,600	22	DEC	DEC	DEC	
- E Europe/E Medit. & Sahelian Africa	10,000-50,000	40,000-65,000	36,000-54,000	7	DEC	DEC	DEC	
- Western Asia/SW Asia & NE Africa	5,000	C	C		DEC	?	DEC	66
<i>Aythya fuligula</i>								
- Northwestern Europe (win)	1,000,000	1,200,000	1,200,000		INC	INC	STA	96
- Central Europe, Black Sea & Med (win)	600,000	700,000	700,000		INC	INC	STA	96
- W Siberia/SW Asia & NE Africa	200,000	200,000	200,000		?	?	?	
<i>Aythya marila marila</i>								
- Northern Europe/Western Europe	310,000	310,000	310,000		?	STA	STA	
- W Siberia/Black Sea & Caspian	100,000-200,000	100,000-200,000	100,000-200,000		?	?	?	
<i>Somateria mollissima mollissima</i>								
- Baltic, Denmark & Netherlands	1,350,000-1,700,000	850,000-1,200,000	760,000	19	STA	DEC	DEC	19
- Norway & Russia	300,000-550,000	360,000-540,000	300,000-550,000	72	STA	STA	STA	
<i>Somateria mollissima borealis</i>								
- Svalbard & Franz Joseph (bre)	40,000-80,000	40,000-80,000	40,000-80,000		STA	STA	STA	
<i>Somateria spectabilis</i>								
- E Greenland, NE Europe & W Siberia	300,000	300,000	300,000		(STA)	STA	STA	
<i>Polysticta stelleri</i>								
- Western Siberia/NE Europe	40,000	30,000-50,000	10,000-15,000	100 48	INC	STA/INC	DEC	100
<i>Clangula hyemalis</i>								
- Iceland & Greenland	150,000	100,000-150,000	100,000-150,000		STA	STA	STA	
- Western Siberia/North Europe	4,600,000	4,600,000	4,600,000		STA	STA	STA	
<i>Melanitta nigra nigra</i>								
- W Siberia & N Eur/W Eur & NW Africa	1,600,000	1,600,000	1,600,000		STA	STA	STA	
<i>Melanitta fusca fusca</i>								
- W Siberia & N Europe/NW Europe	1,000,000	1,000,000	1,000,000		STA	STA	STA	
- Black Sea & Caspian	1,500	1,500	1,500		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Bucephala clangula clangula</i>								
- NW & Central Europe (win)	300,000	400,000	1,000,000-1,300,000	7	INC	INC	STA	96
- NE Europe/Adriatic	75,000	75,000	200,000	7	?	?	?	
- W Siberia & NE Europe/Black Sea	20,000	20,000	60,000	7	?	?	?	
- Western Siberia/Caspian	25,000	25,000	D	70 7	?	?	?	
<i>Mergellus albellus</i>								
- NW & Central Europe (win)	25,000-30,000	40,000	40,000		STA	INC	?	
- NE Eur/Black Sea & E Mediterranean	35,000	35,000	35,000		?	?	?	
- Western Siberia/SW Asia	30,000	30,000	30,000		(DEC)	(DEC)	?	
<i>Mergus serrator serrator</i>								
- NW & Central Europe (win)	145,000	170,000	170,000		STA	INC	?	96
- NE Europe/Black Sea & Med	50,000	50,000	50,000		?	?	?	
- Western Siberia/SW & Cent Asia	<10,000	<10,000	<10,000		?	?	?	
<i>Mergus merganser merganser</i>								
- NW & Central Europe (win)	200,000	250,000	266,100	16 46	STA	INC	DEC	96
- NE Europe/Black Sea	10,000	10,000	10,000		?	?	?	
- Western Siberia/Caspian	20,000	20,000	20,000		?	?	?	
GRUIDAE								
<i>Balearica pavonina pavonina</i>								
- West Africa (Senegal to Chad)	11,500-17,500	15,000	15,000		DEC	DEC	DEC	
<i>Balearica pavonina ceciliae</i>								
- Eastern Africa (Sudan to Uganda)	55,000-60,000	25,000-55,000	28,000-55,000	5	DEC	DEC	DEC	
<i>Balearica regulorum regulorum</i>								
- S Africa (N to Angola & S Zimbabwe)	<10,000	8,000-12,000	7,000-9,000	5	DEC	STA	STA	
<i>Balearica regulorum gibbericeps</i>								
- Eastern Africa (Kenya to Mozambique)	75,000-85,000	50,000-65,000	43,000-55,000	5	DEC	DEC	DEC	
<i>Grus leucogeranus</i>								
- Iran (win)	9	3	4	40	STA	DEC	DEC	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Grus virgo</i>								
- Black Sea (Ukraine)/NE Africa	500	450-510	600-750	7	DEC	DEC	STA	7
- Turkey (bre)	<100	60-90	30-60	7	?	DEC	DEC	
- Kalmykia/NE Africa	30,000-35,000	30,000-35,000	60,000-75,000	7	STA	STA/INC	INC	7
<i>Grus paradisea</i>								
- Extreme Southern Africa	21,000	20,000-21,000	>25,500	55	DEC	STA	STA	
<i>Grus carunculatus</i>								
- Central & Southern Africa	13,000-15,000	8,000	<7,550	5	DEC	DEC	DEC	
<i>Grus grus</i>								
- Northwest Europe/Iberia & Morocco	60,000-70,000	75,000	150,000	56	INC	INC	INC	7
- NE & Central Europe/North Africa	>60,000	70,000	90,000	56	STA/INC	DEC	INC	7
- E Eur/Turkey, Mid. East & NE Africa	35,000	35,000	35,000		DEC	DEC	DEC	
- Turkey & Georgia (bre)	200-500	300-900	200-500	95	DEC	DEC	DEC	
- Western Siberia/South Asia	55,000	70,000	70,000		DEC	?	?	
RALLIDAE								
<i>Sarothrura elegans elegans</i>								
- NE, Eastern & Southern Africa	?	?	?		?	?	(STA)	78
<i>Sarothrura elegans reichenovi</i>								
- S West Africa to Central Africa	?	?	?		?	?	?	
<i>Sarothrura boehmi</i>								
- Central Africa	?	?	A	25	?	DEC	DEC	
<i>Sarothrura ayresi</i>								
- Ethiopia & Southern Africa ⁴	<1,000	700			?	DEC		
Ethiopia			450-650	9			DEC	9
Southern Africa			235	9			DEC	9
<i>Rallus aquaticus aquaticus</i>								
- Europe & North Africa	D	390,000-1,170,000	D	7	STA	STA	DEC	7
<i>Rallus aquaticus korejewi</i>								
- Western Siberia/Southwest Asia	?	?	?		?	?	?	
<i>Rallus caerulescens</i>								
- Southern & Eastern Africa	?	?	?		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Crecopsis egregia</i>								
- Subsaharan Africa	?	?	?		?	?	?	
<i>Crex crex</i>								
- Europe & West Asia/Subsaharan Africa	D	3.4-6.0 million	E	7	DEC	DEC	DEC	
<i>Amaurornis flavirostris</i>								
- Subsaharan Africa	E	E	E		?	?	?	
<i>Porzana parva parva</i>								
- Western Eurasia/Africa	C/D	D	D		DEC	DEC	DEC	
<i>Porzana pusilla intermedia</i>								
- Europe (bre)	B	10,000-20,000	2,000-10,000	7	DEC	STA/DEC	?	7
<i>Porzana porzana</i>								
- Europe/Africa	D	D	D		DEC	DEC	FLU	7
<i>Aenigmatolimnas marginalis</i>								
- Subsaharan Africa	?	?	A/B	21	(DEC)	DEC	DEC	
<i>Porphyrio alleni</i>								
- Subsaharan Africa	?	?	C/D	21	?	?	?	
<i>Gallinula chloropus chloropus</i>								
- Europe & North Africa	E	2.6-4.5 million	>2.7-5.1 million	7	STA	STA	STA	
- West & Southwest Asia	D	D	D		?	?	?	
<i>Gallinula angulata</i>								
- Subsaharan Africa	?	C/D	C/D		?	?	?	
<i>Fulica cristata</i>								
- Subsaharan Africa	D/E	D/E	D	25	?	?	?	
- Spain & Morocco	A	5,000-10,000	5,000	29	?	?	DEC	29
<i>Fulica atra atra</i>								
- Northwest Europe (win)	1,500,000	1,750,000	1,750		STA	INC	STA	96
- Black Sea & Mediterranean (win)	2,500,000	2,500,000	2,500,000		(STA)	(INC)	?	96
- Southwest Asia (win)	2,000,000	2,000,000	2,000,000		?	?	?	
DROMADIDAE								
<i>Dromas ardeola</i>								
- NW Indian Ocean, Red Sea & Gulf	43,000	43,000	60,000-80,000	21 73	?	(STA)	(STA)	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
HAEMATOPODIDAE								
<i>Haematopus ostralegus ostralegus</i>								
- Europe/S & W Europe & NW Africa	1,000,000	1,020,000	1,020,000		INC	INC	DEC	96 75
<i>Haematopus ostralegus longipes</i>								
- SE Eur & W Asia/SW Asia & NE Africa	25,000	100,000-200,000	100,000-200,000		?	(STA)	(DEC)	7
<i>Haematopus moquini</i>								
- Coastal Southern Africa	4,780	4,800	5,000-6,000	21	?	STA	INC	21
RECURVIROSTRIDAE								
<i>Himantopus himantopus himantopus</i>								
- Sub-Saharan Africa (excluding south)	?	D	100,000-200,000	22	?	?	?	
- Southern Africa (' <i>meridionalis</i> ')	-	15,000-30,000	15,000-30,000		-	INC	INC	
- SW Europe & NW Africa/W Africa	40,000	71,000-82,000	71,000-82,000		STA	STA	STA	
- Central Europe & E Medit./N-C Africa	30,000-60,000	23,000-44,000	40,000-60,000	80	STA	?	STA?	80
- W, C & SW Asia/SW Asia & NE Africa	B	20,000-50,000	30,000-70,000	80 7	?	?	?	
<i>Recurvirostra avosetta</i>								
- Southern Africa	10,000-20,000	19,300	19,300		INC	(INC)	INC?	76
- Eastern Africa	?	C	C		?	?	?	
- Western Europe & NW Africa (bre)	67,000	73,000	73,000		INC	STA	STA	
- SE Europe, Black Sea & Turkey (bre)	C	47,000	47,000		(DEC)	STA/DEC	STA/DEC?	76
- W & SW Asia/Eastern Africa	B	B	B		?	(STA)	STA?	
BURHINIDAE								
<i>Burhinus senegalensis senegalensis</i>								
- West Africa	?	B	B		?	?	?	
<i>Burhinus senegalensis inornatus</i>								
- Northeast & Eastern Africa	?	B	B		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
GLAREOLIDAE								
<i>Pluvianus aegyptius aegyptius</i>								
- West & Eastern Africa	?	B			?	?		
West Africa			20,000-50,000	21			STA	21
East Africa			A/B	21			DEC	21
Lower Congo basin			A	21			?	21
<i>Glareola pratincola pratincola</i>								
- W Europe & NW Africa/West Africa	16,000-20,000	18,000-19,500	18,000-19,500		?	STA	STA	
- Black Sea & E Med/E Sahel zone	B	16,000-31,000	16,000-32,000	80	DEC	DEC	DEC	
- SW Asia/SW Asia & NE Africa	B/C	B/C	B/C		?	?	?	
<i>Glareola nordmanni</i>								
- SE Europe & W Asia/Southern Africa	D	29,000-45,000	29,000-45,000		DEC	DEC	DEC	
<i>Glareola ocularis</i>								
- Madagascar/East Africa	?	A/B	5,000-10,000	21	?	?	DEC	21
<i>Glareola nuchalis nuchalis</i>								
- Eastern & Central Africa	?	C	C		DEC	?	?	
<i>Glareola nuchalis liberiae</i>								
- West Africa	?	B/C	>100,000	27	?	?	?	
<i>Glareola cinerea cinerea</i>								
- SE West Africa & Central Africa	?	B/C	B	21	?	?	?	
CHARADRIIDAE								
<i>Pluvialis apricaria apricaria</i>								
- Britain, Ireland, DK, Ger & Baltic (bre)	70,000	69,000	140,000-210,000	80 7	DEC	DEC	DEC	
<i>Pluvialis apricaria altifrons</i>								
- Iceland & Faroes/E Atlantic coast	750,000	930,000	930,000		(STA)	(STA)	(STA)	
- N Europe/W Europe & NW Africa	1,000,000	645,000-954,000	500,000-1,000,000		STA	STA	STA	
- Northern Siberia/Caspian & Asia Minor	?	?	?		?	?	?	
<i>Pluvialis fulva</i>								
- N-C Siberia/S & SW Asia, NE Africa	C/D	50,000-100,000	50,000-100,000		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Pluvialis squatarola</i>								
- W Siberia & Can/W Europe & W Africa	168,000	247,000	247,000		INC	INC	DEC	96 75
- C & E Siberia/SW Asia, E & S Africa	50,000	90,000	90,000		?	?	?	
<i>Charadrius hiaticula hiaticula</i>								
- N Europe/Europe & North Africa	47,500	73,000	73,000		INC	INC	DEC	96 75
<i>Charadrius hiaticula psammodroma</i>								
- Can, Greenl. & Iceland/W & S Africa	195,000	190,000	190,000		?	(DEC)	DEC	76
<i>Charadrius hiaticula tundrae</i>								
- NE Eur & Sib/SW Asia, E & S Africa	200,000	145,000-280,000	D	75	?	?	?	
<i>Charadrius dubius curonicus</i>								
- Europe & NW Africa/W Africa	D	180,000-290,000	200,000-300,000	7	STA	(STA)	STA?	76
- West & SW Asia/Eastern Africa	?	?	?		?	?	?	
<i>Charadrius pecuarius pecuarius</i>								
- Southern & E & NE Africa	?	50,000-100,000	100,000-400,000	21	?	?	?	
- West Africa	-	10,000-20,000	20,000-50,000	21	-	?	?	
<i>Charadrius tricollaris tricollaris</i>								
- Southern & Eastern Africa	?	40,000-100,000	70,000-130,000	21	?	INC	INC	
<i>Charadrius forbesi</i>								
- West & Central Africa	?	B/C	B/C		?	?	?	
<i>Charadrius pallidus pallidus</i>								
- Southern Africa	6,000-7,000	11,200	11,200		STA	STA	STA	
<i>Charadrius pallidus venustus</i>								
- Eastern Africa	B	4,000-5,000	4,000-5,000		?	?	?	
<i>Charadrius alexandrinus alexandrinus</i>								
- West Europe & W Med/West Africa	67,000	62,000-70,000	62,000-70,000		DEC	DEC	DEC	
- Black Sea & E Medit./E Sahel	C	32,000-49,000	32,000-49,000		DEC	(DEC)	DEC	7
- SW & Cen Asia/SW Asia & NE Africa	C/D	C	C		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Charadrius marginatus mechowii</i>								
Southern & Eastern Africa	?	25,000-35,000			?	?		
Inland Eastern to Southern Africa			10,000-15,000	64			?	
Coastal E Africa			15,000-25,000	21			?	
West to West-central Africa	?	10,000-15,000	10,000-15,000		?	?	?	
<i>Charadrius mongolus pamirensis</i>								
- West-central Asia/SW Asia & E Africa	>30,000	>30,000	30,000-50,000	21	?	?	?	
<i>Charadrius leschenaultii columbinus</i>								
- Turkey & SW Asia/E. Med & Red Sea	A	A	A		?	?	?	
<i>Charadrius leschenaultii crassirostris</i>								
- Caspian & SW Asia/Arabia & NE Africa	65,000	C	C		?	?	?	
<i>Charadrius leschenaultii leschenaultii</i>								
- Central Asia/Eastern & Southern Africa	-	B	25,000-50,000	21	-	?	?	
<i>Charadrius asiaticus</i>								
Central Asia/ E & southern Africa	B/C	B/C	40,000-55,000	21	DEC	(DEC)	(DEC)	
<i>Eudromias morinellus</i>								
- Europe/NW Africa	D	39,000-110,000	40,000-120,000	7	DEC	STA/DEC	(STA/DEC)	76
- Asia/Middle East	B/C	B/C	B/C		?	?	?	
<i>Vanellus vanellus</i>								
- Europe/Europe & N Africa	7,000,000	2.8-4.0 million	5.1-8.4 million	80 7	DEC	DEC	DEC	
- Western Asia/SW Asia	C/D	1.6-2.9 million	C/D	66	?	?	?	
<i>Vanellus spinosus</i>								
- Black Sea & Mediterranean (bre)	C	C	C		INC	INC	INC	
<i>Vanellus albiceps</i>								
- West & Central Africa	?	10,000-20,000	30,000-70,000	21	?	?	STA	22
<i>Vanellus senegallus senegallus</i>								
- West Africa	?	C	25,000-60,000	21	?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Vanellus senegallus solitaneus</i>								
- Southwest Africa	?	C	B/C	21	?	?	?	
<i>Vanellus senegallus lateralis</i>								
- Eastern & Southeastern Africa	?	C	C		?	?	STA	35
<i>Vanellus lugubris</i>								
- Southern West Africa	-	5,000-20,000	5,000-20,000		-	(DEC)	?	
- Central & Eastern Africa	?	20,000-50,000	20,000-50,000		?	DEC	?	
<i>Vanellus melanopterus minor</i>								
- Southern Africa	?	2,000-3,000	2,000-3,000		(DEC)	DEC	DEC	
<i>Vanellus coronatus coronatus</i>								
- Eastern & Southern Africa	?	200,000-400,000	400,000-900,000	21	?	(INC)	?	
- Central Africa	?	?	A/B	21	?	?	?	
<i>Vanellus coronatus xerophilus</i>								
- Southwest Africa	?	?	B/C	21	?	?	INC	35
<i>Vanellus superciliosus</i>								
- West & Central Africa	?	A/B	A/B		?	?	?	
<i>Vanellus gregarius</i>								
- SE Europe & Western Asia/NE Africa	A	400-1,200	400-1,200		DEC	DEC	DEC	
- Central Asian Republics/NW India	<1,000	200-600	200-600		DEC	DEC	DEC	
<i>Vanellus leucurus</i>								
- SW Asia/SW Asia & NE Africa	B/C	B	B		(INC)	(DEC)	(DEC)	
- Central Asian Republics/South Asia	B/C	B/C	B/C		?	(INC)	(INC)	
SCOLOPACIDAE								
<i>Scolopax rusticola</i>								
- Europe/S & W Europe & North Africa	>15,000,000	21-25 million	10-25 milion	80	STA	STA	STA	
- Western Siberia/SW Asia (Caspian)	?	?	?		?	?	?	
<i>Gallinago stenura</i>								
- N Siberia/South Asia & Eastern Africa	C/D	C/D	C/D		?	?	?	
<i>Gallinago media</i>								
- Scandinavia/probably West Africa	18,000-51,000	18,000-51,000	18,000-51,000		STA	STA	STA	
- W Siberia & NE Europe/SE Africa	D	D	D		DEC	(DEC)	(DEC)	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Gallinago gallinago gallinago</i>								
- Europe/S & W Europe & NW Africa	>20,000,000	E	>2.5 million	80	DEC	DEC	DEC/ST A	76 75
- W Siberia/SW Asia & Africa	E	>1,500,000	>1,500,000		?	?	?	
<i>Gallinago gallinago faeroensis</i>								
- Iceland, Faroes & N Scotland/Ireland	750,000	570,000	570,000		STA	(STA)	(STA)	
<i>Lymnocyptes minimus</i>								
- N Europe/S & W Europe & W Africa	C/D	D/E	E	45	DEC	STA	STA	
- W Siberia/SW Asia & NE Africa	?	?	?		?	?	?	
<i>Limosa limosa limosa</i>								
- Western Europe/NW & West Africa	350,000	148,000-183,000	162,000-183,000	7	DEC	DEC	DEC	
- Eastern Europe/Central & Eastern Africa	D	93,000-173,000	90,000-165,000	7	DEC	DEC	DEC	
- W-cen. Asia/SW Asia & Eastern Africa	C	C	C		?	?	?	
<i>Limosa limosa islandica</i>								
- Iceland/Western Europe	65,000	35,000	47,000	31	INC	INC	INC	
<i>Limosa lapponica lapponica</i>								
- N Europe/Western Europe	115,000	120,000	120,000		STA/DE C	STA	STA	
<i>Limosa lapponica taymyrensis</i>								
- W Siberia/West & Southwest Africa	700,000	520,000	600,000	75	?	DEC	INC	89
<i>Limosa lapponica taymyrensis</i>								
- Cen Siberia/S & SW Asia & E Africa	C/D	100,000-150,000	100,000-150,000		?	?	?	
<i>Numenius phaeopus phaeopus</i>								
- Northern Europe/West Africa	-	156,000-298,000	190,000-340,000	80 7	-	?	(STA)	7
- West Siberia/Southern & Eastern Africa	?	D	D		?	?	?	
<i>Numenius phaeopus islandicus</i>								
- Iceland, Faroes & Scotland/West Africa	-	610,000	600,000-750,000	80 7	-	STA	(STA)	7
<i>Numenius phaeopus alboaxillaris</i>								
- SW Asia/Eastern Africa	A	A	A		DEC	?	DEC	58

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Numenius tenuirostris</i>								
- Cen Siberia/Mediterranean & SW Asia	50-270	<50	<50		DEC	DEC	DEC	
<i>Numenius arquata arquata</i>								
- Europe/Europe, North & West Africa	348,000	420,000	700,000-1,000,000	80 7	STA/IN C	STA/IN C	DEC	76 96
<i>Numenius arquata orientalis</i>								
- W Siberia/SW Asia, E & S Africa	C	C	C		(DEC)	(DEC)	(DEC)	
<i>Numenius arquata suschkini</i>								
- SE Europe & SW Asia (bre)	-	?	?		-	(DEC)	DEC	76
<i>Tringa erythropus</i>								
- N Europe/S Europe, N & W Africa	75,000-150,000	77,000-131,000	60,000-120,000	80 7	(STA)	(STA)	(STA)	
- W Siberia/SW Asia, NE & East Africa	B/C	B/C	B/C		?	?	?	
<i>Tringa totanus totanus</i>								
- NW Europe/W Europe, NW & W Africa	177,000	222,500	250,000	76 21	DEC	DEC	DEC	
- Central & East Europe/E Med & Africa	D	223,000-464,000	223,000-464,000		DEC	DEC	DEC	
<i>Tringa totanus britannica</i>								
- Britain & Ireland/Britain, Ireland, France	-	124,000-127,000	95,000-135,000	80 7	-	DEC	DEC	
<i>Tringa totanus ussuriensis</i>								
- West Asia/SW Asia, NE & E Africa	?	213,000-326,000	D	75	?	?	?	
<i>Tringa totanus robusta</i>								
- Iceland & Faroes/Western Europe	150,000-300,000	64,500	150,000-400,000	80 7	STA/IN C	STA/IN C	(STA/IN C)	76
<i>Tringa stagnatilis</i>								
- Eastern Europe/West & Central Africa	C/D	21,000-52,000	14,000-40,000	76	?	?	(DEC)	7
- W Asia/SW Asia, E & S Africa	C	C	50,000-100,000	21	?	?	?	
<i>Tringa nebularia</i>								
- N Europe/SW Europe, NW & W Africa	D	234,000-395,000	190,000-270,000	80 7	STA	STA	STA	
- Western Siberia/SW Asia, E & S Africa	C/D	D	D		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Tringa ochropus</i>								
- N Europe/S & W Europe, West Africa	D/E	1.0-1.9 million	1.0-2.4 million	80 7	STA/IN C	STA/IN C	STA	76
- W Siberia/SW Asia, NE & E Africa	?	D/E	D/E		?	?	?	
<i>Tringa glareola</i>								
- NW Europe/West Africa	E	855,000-1,220,000	900,000-1,200,000	80 7	DEC	DEC	STA	7
- NE Europe & W Siberia/E & S Africa	D/E	>2,000,000	>2,000,000		?	(STA)	(STA)	
<i>Xenus cinereus</i>								
- NE Eur & W Sib/SW Asia, E & S Africa	44,000	D	D		(STA)	(STA)	(STA)	
<i>Actitis hypoleucos</i>								
- W & Central Europe/West Africa	E	1.4-2.0 million	1.5-2.0 million	80 7	STA	STA	(DEC)	7
- E Europe & W Siberia/C, E & S Africa	(E)	E	E		?	(STA)	(STA)	
<i>Arenaria interpres interpres</i>								
- NE Can & Greenl/W Eur & NW Africa	>80,000	100,000-200,000	100,000-200,000		(INC)	INC	DEC	96 75
- Northern Europe/West Africa	50,000-100,000	46,000-119,000	45,000-120,000	80 7	STA	STA	DEC	75
- W & Cen Siberia/SW Asia, E & S Africa	C	100,000	100,000		?	?	?	
<i>Calidris tenuirostris</i>								
- E Siberia/SW Asia & W South Asia	A	2,000-5,000	2,000-5,000		?	?	?	
<i>Calidris canutus canutus</i>								
- N Siberia/Western & Southern Africa	260,000	340,000	500,000	21 89	DEC	DEC	?	75
<i>Calidris canutus islandica</i>								
- NE Canada & Greenland/West Europe	400,000	450,000	450,000		(STA)	DEC	DEC	
<i>Calidris alba</i>								
- E Atlantic, W & S Africa (win)	123,000	123,000	123,000		STA	STA/IN C	STA/IN C	
- SW Asia, E & S Africa (win)	120,000	140,000	150,000	21 76	?	(STA)	(STA)	
<i>Calidris minuta</i>								
- N Europe/Europe, North & West Africa	211,000	200,000	>200,000	22	STA	(DEC)	(DEC)	
- Western Siberia/SW Asia, E & S Africa	1,000,000	1,000,000	1,000,000		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Calidris temminckii</i>								
- Fennoscandia/North & West Africa	?	39,000-80,000	39,000-80,000		?	?	?	
- NE Eur & W Sib/SW Asia & E Africa	?	(E)	D/E	22	?	?	?	
<i>Calidris maritima maritima</i>								
- North & W Europe (excl. Iceland) (win)	50,500	50,000-100,000	50,000-100,000		STA	STA	STA	
<i>Calidris alpina alpina</i>								
- NE Eur & NW Sib/W Eur & NW Africa	1,373,000	1,330,000	1,330,000		DEC	STA	STA	
<i>Calidris alpina centralis</i>								
- Central Siberia/SW Asia & NE Africa	150,000	300,000	500,000	75	?	?	?	
<i>Calidris alpina schinzii</i>								
- Iceland & Greenland/NW and W Africa	800,000	940,000-960,000	940,000-960,000		STA	STA	STA	
- U.K. & Ireland/SW Eur & NW Africa	33,000-36,000	23,000-26,000	23,000-26,000		DEC	DEC	DEC	
- Baltic/SW Europe & NW Africa	-	3,600-4,700	3,300-4,100	80 7	-	DEC	DEC	
<i>Calidris alpina arctica</i>								
- NE Greenland/West Africa	15,000	21,000-45,000	21,000-45,000		STA	(STA)	(STA)	
<i>Calidris ferruginea</i>								
- Western Siberia/West Africa	436,000	740,000	1,000,000	89	STA	INC	INC	
- Central Siberia/SW Asia, E & S Africa	310,000	330,000	>400,000	21	STA	STA	STA	
<i>Limicola falcinellus falcinellus</i>								
- Northern Europe/SW Asia & Africa	40,000-60,000	61,000-64,000	61,000-64,000		(DEC)	?	DEC	7
<i>Philomachus pugnax</i>								
- N Europe & W Siberia/West Africa	E	E	1.0 million-1.5 million	85	DEC	DEC	DEC	
- Northern Siberia/SW Asia, E & S Africa	D/E	E	E		?	(DEC)	(DEC)	
<i>Phalaropus lobatus</i>								
- Western Eurasia/Arabian Sea	D/E	E	E		(STA)	?	?	
<i>Phalaropus fulicarius</i>								
- Can & Greenland/Atlantic coast Africa	?	920,000	1,000,000	59	?	?	DEC	59

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
LARIDAE								
<i>Larus leucophthalmus</i>								
- Red Sea & nearby coasts	20,000	20,000	37,000-44,000	67 22	STA	STA	STA	
<i>Larus hemprichii</i>								
- Red Sea, Gulf, Arabia & Eastern Africa	D	150,000-300,000	150,000-300,000		?	?	(INC)	67
<i>Larus canus canus</i>								
- NW & C Europe/Atlantic & Medit.	1,600,000	1.3-2.1 million	1.2-2.25 million	7	DEC	DEC	(DEC)	7
<i>Larus canus heinei</i>								
- NE Eur & W Sib/Black Sea & Caspian	C	D	E	7	?	(INC)	(INC)	7
<i>Larus audouinii</i>								
- Mediterranean/N & W coasts of Africa	40,000	57,600	57,600		INC	INC	INC	
<i>Larus marinus</i>								
- North & Western Europe	480,000	420,000-510,000	330,000-540,000	7	INC	STA	INC	7
<i>Larus dominicanus vetula</i>								
- Coastal Southern Africa	?	69,000	70,000	13	?	INC	INC	
<i>Larus hyperboreus hyperboreus</i>								
- North Atlantic	200,000	135,000-360,000	135,000-360,000		STA	STA	STA	
<i>Larus hyperboreus leuceteres</i>								
Canada Greenland & Iceland (bre)			D				?	
<i>Larus glaucooides glaucooides</i>								
- Greenland/Iceland & NW Europe	C/D	90,000-300,000	90,000-300,000		STA	STA	STA	
<i>Larus argentatus argentatus</i>								
- North & Northwest Europe	1,400,000	1.1-1.5 million	1.7-3.6 million	7	INC	INC	INC	
<i>Larus argentatus argenteus</i>								
- Iceland & Western Europe	1,300,000	1,090,000	560,000-620,000	7	?	(STA)	DEC	7
<i>Larus heuglini</i>								
- NE Eur & W Sib/W Asia & NE Africa	?	?	?		?	?	?	
<i>Larus (heuglini) barabensis</i>								
- Southwest Siberia/Southwest Asia	?	?	?		?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Larus armenicus</i>								
- Armenia, E Turkey & NW Iran	45,000-60,000	69,000-75,000	69,000-75,000		STA	?	?	
<i>Larus cachinnans cachinnans</i>								
- Black S. & W Asia/SW Asia, NE Africa	C/D	D	D		?	(INC)	INC	7
<i>Larus cachinnans michahellis</i>								
- Mediterranean, Iberia & Morocco	350,000	475,000-585,000	630,000-768,000	94	INC	INC	INC	
<i>Larus fuscus fuscus</i>								
- NE Eur/Black Sea, SW Asia & E Africa	200,000-300,000	156,000-228,000	56,000	34	?	DEC	DEC	34
<i>Larus fuscus graellsii</i>								
- Western Europe/Med & West Africa	400,000-500,000	525,000	530,000-570,000	7	INC	INC	INC	7
<i>Larus fuscus intermedius</i>								
S Scandinavia, Netherlands, Ebro Delta, Spain			325,000-440,000	7 34			INC	7
<i>Larus ichthyæetus</i>								
- Black Sea & Caspian/SW Asia	70,000-120,000	72,000-120,000	D	7	INC	INC	INC	
<i>Larus cirrocephalus poiocephalus</i>								
- West Africa	?	30,000	30,000		?	?	?	
- Central & Eastern Africa	?	D			?	?		
Coastal Southern Africa			200,000-400,000	21			?	
- Southern Africa (excluding Madagascar)	?	(B)			?	INC		
<i>Larus hartlaubii</i>								
- Coastal Southwest Africa	25,000	32,000-33,000	30,000	81	?	STA/INC	INC	81
<i>Larus ridibundus</i>								
- W Europe/W Europe, W Med, W Africa	>5,000,000	5.6-7.3 million	3.7-4.8 million	7	(STA)	INC	DEC	7
- E Europe/Black Sea & E Mediterranean	D	1.3-1.7 million	770,000-1,800,000	7	(STA)	INC	?	7
- West Asia/SW Asia & NE Africa	250,000	250,000	250,000		?	?	?	
<i>Larus genei</i>								
- West Africa (bre)	20,000	22,500	22,500		INC	INC	STA	21
- Black Sea & Mediterranean (bre)	120,000-240,000	123,000-237,000	140,000-205,000	7	(INC)	INC	STA	7
- W, SW & S Asia (bre)	150,000	150,000	150,000		INC	INC	INC	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Larus melanocephalus</i>								
- W Europe, Med & NW Africa	D	570,000-1,110,000	360,000-960,000	7	INC	INC	STA	7
<i>Larus minutus</i>								
- Cen & E Europe/SW Europe & W Med.	60,000-90,000	66,000-102,000	72,000-174,000	7	STA/INC	(STA)	INC	7
- W Asia/E Med, Black Sea & Caspian	(C)	C	C		?	?	?	
<i>Xema sabini sabini</i>								
- Canada & Greenland/SE Atlantic	?	?	300,000-600,000	49	?	?	?	
<i>Sterna nilotica nilotica</i>								
- Western Europe/West Africa	12,000	9,500-10,800	14,000-21,000	7 21	DEC	DEC	STA	7
- Black Sea & E Medit./Eastern Africa	?	14,000-39,000	24,000-52,000	7	DEC	DEC	DEC	
- West & Central Asia/SW Asia	B	B	B		?	?	?	
<i>Sterna caspia caspia</i>								
- Southern Africa (bre)	1,500	5,000	2,000	22 84	STA	(INC)	STA	81
- West Africa (bre)	15,000	40,500	45,000-60,000	21 92	(DEC)	INC	STA	21
- Europe (bre)	5,000-7,000	5,400-7,800	8,000-11,000	7	DEC	DEC	INC	7
- Caspian (bre)	10,000	9,000-16,500	9,000-16,500		DEC	(DEC)	?	
<i>Sterna maxima albidorsalis</i>								
- West Africa (bre)	75,000	129,000	225,000	93	STA	STA/INC	STA/INC	
<i>Sterna bengalensis bengalensis</i>								
- Persian Gulf/Southern Asia	150,000-180,000	150,000-180,000	150,000-180,000		?	?	?	
<i>Sterna bengalensis par</i>								
- Red Sea/Eastern Africa	C	C	40,000-47,000	67 21	?	?	STA	21
- S Mediterranean/NW & W Africa coasts	4,000	4,000	4,000		STA	STA	STA	
<i>Sterna bergii bergii</i>								
- Southern Africa (Angola – Mozambique)	20,000	20,000	20,000		?	(STA)	(STA)	
<i>Sterna bergii enigma</i>								
- Madagascar & Mozambique/S Africa	-	8,000-10,000	7,500-10,000	21	-	?	?	
<i>Sterna bergii thalassinus</i>								
- Eastern Africa & Seychelles	1,200	2,550-4,500	1,300-1,700	21	?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Sterna bergii velox</i>								
- Red Sea & NE Africa	C	C	C		?	?	?	
<i>Sterna sandvicensis sandvicensis</i>								
- Western Europe/West Africa	150,000	159,000-171,000	166,000-171,000	7	INC	INC	STA	7
- Black Sea & Mediterranean (bre)	130,000	44,000-73,000	61,000-197,000	7	(DEC)	DEC	FLU	7
- West & Central Asia/SW & South Asia	110,000	110,000	110,000		?	?	?	
<i>Sterna dougallii dougallii</i>								
- Southern Africa	400	750	750-780	51	DEC	INC		
- East Africa	38,000	26,000	25,500	12	?	?	?	
- Europe (bre)	5,000	4,800-5,400	5,400-5,700	7	DEC	DEC	INC	7
<i>Sterna dougallii arideensis</i>								
- Madagascar, Seychelles & Mascarenes	3,600	10,000-15,000	12,000-15,000	21	?	(DEC)	DEC	21
<i>Sterna dougallii bangsi</i>								
- North Arabian Sea (Oman)	?	<600	<600		?	?	?	
<i>Sterna vittata vittata</i>								
- Subantarctic islands/S Africa	3,000-6,000	3,000-6,000	>6,700	81	?	?	?	
<i>Sterna vittata tristanensis</i>								
- Tristan da Cunha & Gough/S Africa	2,500	2,500	2,400-4,500	83	?	?	?	
<i>Sterna hirundo hirundo</i>								
- Southern & Western Europe (bre)	180,000	170,000-200,000	170,000-210,000	7	(STA)	(STA)	STA	7
- Northern & Eastern Europe (bre)	600,000	460,000-820,000	630,000-1,500,000	7	(STA)	(STA)	STA	7
- Western Asia (bre)	C/D	C/D	C/D		?	?	?	
<i>Sterna paradisaea</i>								
- Western Eurasia (bre)	E	1.3-2.3 million	E	7	(STA)	STA/DEC	?	
<i>Sterna albifrons albifrons</i>								
- Eastern Atlantic (bre)	34,000	31,000-37,500	42,500-55,500	7	STA/INC	STA	DEC	7
- Black Sea & East Mediterranean (bre)	70,000-120,000	63,500-127,000	63,500-112,500	7	DEC	DEC	DEC	
- Caspian (bre)	B	B	B		?	?	?	
<i>Sterna albifrons guineae</i>								
- West Africa (bre)	?	?	2,000-3,000	21	?	?	?	

Population	CSR 1999	CSR 2002	Proposed revision 2005	Source 2005 revision	Trend 1999	Trend 2002	Proposed Trend 2005	Source 2005 trend
<i>Sterna saundersi</i>								
- W South Asia, Red Sea, Gulf & E Africa	40,000	40,000	40,000		?	?	?	
<i>Sterna balaenarum</i>								
- Namibia & S Africa/Atlantic to Ghana	13,500	13,500	14,000	74 81	(DEC)	(INC)	STA	21
<i>Sterna repressa</i>								
- W S Asia, Red Sea, Gulf & E Africa	600,000	600,000	600,000		DEC	DEC	DEC	
<i>Chlidonias hybridus hybridus</i>								
- Western Europe & NW Africa (bre)	20,000-30,000	21,500-31,000	16,300-39,800	7 42	DEC	DEC	STA	7
- Black Sea & E Mediterranean (bre)	50,000-80,000	80,000-120,000	98,000-108,000	7	STA/IN C	STA/IN C	STA	7
- Caspian (bre)	C	C	C		?	?	?	
<i>Chlidonias hybridus sclateri</i>								
- Eastern Africa (Kenya & Tanzania)	?	A	10,000-15,000	21	?	?	?	
- Southern Africa (to Malawi & Zambia)	?	B	5,000-15,000	21	?	?	?	
<i>Chlidonias leucopterus</i>								
- E Europe & W Asia/Africa	200,000-250,000	3,000,000	3,000,000		DEC	?	(STA)	7
<i>Chlidonias niger niger</i>								
- Europe & Asia/Atlantic coast of Africa	200,000	200,000-350,000	500,000-1,000,000	7	DEC	STA/DEC	DEC	7
RYNCHOPIDAE								
<i>Rynchops flavirostris</i>								
- Coastal West Africa	?	<5,000			?	DEC		
- Central Africa	?	B			?	DEC		
Central & coastal West Africa			7,000-13,000	21			DEC	21
- Eastern & Southern Africa	?	B	8,000-12,000	21	?	DEC	DEC	

Sources for Table 1.

1	Azafzaf, H., and Paul Isenmann. In litt. June-August, 2002.
2	Baker, N. 2003. A reassessment of the northern population of the Cape Teal <i>Anas capensis</i> . <i>Scopus</i> 23: 29-43.
3	Béchet, A. & Germain, C. (2004) Greater Flamingo breeding report for the Mediterranean and West Africa, 2004. Station Biologique Tour du Valat, Arles, France.
4	Béchet, A. 2005. Estimate of the Greater Flamingo <i>Phoenicopterus roseus</i> population size in the Western Palearctic in 2005. Unpublished synthesis prepared for the Flamingo Specialist Group and WPE4.
5	Beilfus R., Dodman, T. and Urban, E. (2005) Status of Cranes in Africa. 2004 The Ostrich.
6	van den Berg, A. 2002. WP Reports. <i>Dutch Birding</i> 24: 234-235.
7	BirdLife International (2004). Birds in Europe, population estimates, trends and conservation status. Cambridge, U.K: BirdLife International. (BirdLife Conservation Series No. 12).
8	<u>BirdLife International (2005) Threatened birds of the world 2005. Species factsheets available at www.birdlife.org</u>
9	BirdLife International. (2000). Threatened Birds of the World. Barcelona and Cambridge, UK: Lynx Edicions and BirdLife international
10	Brown, L.H., Urban, E.K. and Newman, K. 1982. The Birds of Africa. Volume I. Academic Press, London and New York.
11	Clausen, P., in litt. 2005.
12	Cooper, J., Williams, A.J. and Britton, P.L. 1984. Distribution, population sizes and conservation of breeding seabirds in the Afrotropical region. IUCN Technical Publication 2, 403-417.
13	Crawford, R. in prep. 2002. Species texts for: Cape Cormorant, <i>Phalacrocorax capensis</i> , Crowned Cormorant, <i>Phalacrocorax coronatus</i> , Great White Pelican, <i>Pelecanus onocrotalus</i> , Kelp Gull, <i>Larus dominicanus</i> . In: Hockey, P.A.R., et al. in prep. , Roberts' Birds of Southern Africa, 7th edition.
14	Crivelli, A.J. and Schreiber, R.W. 1984. Status of the Pelecanidae. <i>Biol. Cons.</i> 30:147-156.
15	Crivelli, A.J. Catsadorakis, G. Hatzilacou, D., Hulea, D., Malakou, M., Marinov, M., Michev, T., Nazirides, T., Peja, N., Sarigul, G. & Siki, M. 2000. Status and population development of Great White Pelican <i>Pelecanus onocrotalus</i> and Dalmatian Pelican <i>P. crispus</i> breeding in the Palearctic. Pp 38-46 In: Yesou, P. & Sultana, J. (eds.) Proceedings of the 5th Medmaravis Symposium, Gozo, Malta. Environment Protection Department, Malta.
16	Delany, S.N., Reyes, C. Hubert, E., Pihl, S., Rees, E., Haanstra, L., & van Strien, A. 1999. Results from the International Waterbird Census in the Western Palearctic and Southwest Asia 1995 and 1996.
17	Denny, M.J.H., Clausen, P., Percival, S.M., Anderson, G.Q.A., Koffijberg, K. & Robinson, J.A. (2004): Light-bellied Brent Goose <i>Branta bernicla hrota</i> [East Atlantic population] in Svalbard, Greenland, Franz Josef Land, Norway, Denmark, The Netherlands and Britain 1960/61 - 2000/01 . The Wildfowl and Wetlands Trust/Joint Nature Conservation Committee, Slimbridge. - <i>Waterbird Review Series</i> : 45 pp.
18	Dereliev, S., in litt. 2005
19	Desholm, M., Christensen, T.K., Scheiffarth, G., Hario, M., Andersson, Å., Ens, B., Camphuysen, C.J., Nilsson, L., Waltho, C.M., Lorentsen, S-H., Kuresoo, A., Kats, R.K.H., Fleet, D.M. & Fox, A.D. 2002. Status of the Baltic/Wadden Sea population of the Common Eider <i>Somateria m. mollissima</i> . <i>Wildfowl</i> 53: 167-203.
20	Diawara, Y., Arnaud, A., Araujo, A. & Béchet, A. In review, 2005. Nouvelles données sur la reproduction et l'hivernage des flamants roses en Mauritanie et confirmation d'échanges avec les populations méditerranéennes. <i>Ostrich</i> (submitted).
21	Dodman, T. 2002. Waterbird Population Estimates in Africa. Unpublished report to Wetlands International.
22	Dodman, T. In press, 2005. Status, estimates and trends of waterbird populations in Africa. Wetlands International, Dakar.
23	Dodman, T., in litt. to SR.
24	Ebbinge, B.S. 2004. Onderzoek naar het broedsucces van Zwartbuikrotganzen [Unravelling the breeding success of Dark-bellied Brent Geese]. <i>Limosa</i> 77: 71-78.

25	Fishpool, L.D.C. & Evans, M.I.,(Eds). 2001. Important Bird Areas in Africa and associated islands: Priority sites for conservation. Newbury and Cambridge, UK: Pisces Publications and BirdLife International (BirdLife Conservation Series No. 11).
26	Fox, A.D. & Francis, I. 2004. Report of the 2003/2004 Census of Greenland White-fronted Geese in Britain. Greenland White-fronted Goose Study, Kalø, Denmark
27	Gatter, W. 1997. Birds of Liberia. Pica Press, Sussex.
28	Gilissen, N., Haanstra, L. & Delany, S. Boere, G. & Hagemeyer, W. 2002. Numbers and Distribution of Wintering Waterbirds in the Western Palearctic and Southwest Asia in 1997, 1998 and 1999 - Results from the International Waterbird Census. Wetlands International Global Series No 11.
29	Green in litt. 2003.
30	Green, A.J. in litt. 2002.
31	Gunnarsson et al. (2005) (Bird Study 52: 145-152).
32	Hafner, H. 2000. Herons in The Mediterranean. Pp 32-54 in: Heron Conservation (James.A. Kushlan & Heinz Hafner, Eds.). Academic Press, London.
33	Harebottle, D., in litt. 2002.
34	Hario, M., in litt. 2005.
35	Harrison, J.A., Allan, D.G., Underhill, L.G., Herremans, M., Tree, A.J., Parker, V. & Brown, C.J. (1997). The Atlas of Southern African Birds. Vol.1. BirdLife South Africa, Johannesburg. 785 pp.
36	Hockey P.A.R., Dean, W.R.J. & Ryan, P.G. (eds). 2005. Roberts - Birds of Southern Africa, 7th ed. The Trustees of the John Voelcker Bird Book Fund, Cape Town.
37	del Hoyo, J., Elliot, A. and Sargatal, J. (eds). 1996. Handbook of the Birds of the World. Volume 3: Hoatzin to Auks. Lynx Edicions, Barcelona.
38	del Hoyo, J., Elliott A. and Sargatal, J. (eds). 1992. Handbook of the Birds of the World. Volume 1: Ostrich to Ducks. Lynx Edicions, Barcelona.
39	Hughes, B., Unpublished r on the Threatened Waterfowl Specialist Group forum, 17 January 2005
40	Ilyashenko, E. 2002. Siberian Crane Flyway Newsletter. CMS/ICF/CWGE
41	Irish Brent Goose Research Group data.
42	Isenmann, P. & Moali, A. 2000. Oiseaux d'Algérie / Birds of Algeria. SEOF, Paris.
43	Jennings, M.C. 2003. ABBA Survey 31: Bir Ali and Red Sea Islands, Yemen. <i>Phoenix</i> 19:11-20.
44	Jennings, M.C. in prep. Atlas of Breeding Birds of Arabia.
45	Kalchreuter, H. 2003. On the population status of the Jack Snipe (<i>Lymnocyptes minimus</i>). <i>Game & Wildlife Science</i> 20 (3): 175-193
46	Kershaw, M. & P.A. Cranswick. 2002 in press. Numbers of wildfowl and selected waterbirds wintering in Great Britain. <i>Biological Conservation</i> .
47	Koffijberg, K., in litt. 2002.
48	Krasnov, Y.V., Ström, H., Gavriilo, M.V., and Shavykin, A.A.. 2004. Seabirds wintering in Polynyas along the Terskiy Coast of the White Sea and along East Murman Coast. <i>Faunistics</i> 31: 51-57.
49	Kushlan, James, A., Melanie J. Steinkamp, Katharine Parsons, Jack Capp, Martin Acosta Cruz, Malcolm Coulter, Ian Davidson, Loney Dickson, Naomi Edelson, Richard Elliot, R. Michael Erwin, Scott Hatch, Stephen Kress, Robert Milko, Steve Miller, Kyra Mills, Richard Paul, Roberto Phillips, Jorge E. Saliva, Bill Sydeman, John Trapp, Jennifer Wheeler, and Kent Wohl. 2002. North American Waterbird Conservation Plan, Version 1. Waterbird Conservation for the Americas, Washington, DC, USA.
50	Li, David & Taej Mundkur, 2002. Status Overview and Recommendations for the White-headed Duck Central Asian Population. in press.
51	Louette, M. 2004. Oiseaux. In: Louette, M., D. Meirte and R. Jocque (eds). La faune terrestre de l'archipel des Comores. <i>Studies in Afrotropical Zoology (MRAC, Tervuren)</i> . 293: 89-196.
52	Madsen, J., Cracknell, G. & Fox, A.D. (eds). (1999). Goose populations of the Western Palearctic: A review of status and distribution. Wetlands International Publication No.48. Wetlands International, Wageningen, The Netherlands; National Environmental Research Institute, Ronde, Denmark. 344 pp.
53	Madsen, J., Kuijken, E., Cottaar, F., unpublished data.

54	Marion, L. Ulenaers, P. & van Vessem, J.. 2000. Herons in Europe. Pp 1-31 in: Heron Conservation (James.A. Kushlan & Heinz Hafner, Eds.). Academic Press, London.
55	McCann et al. 2005. Conservation Priorities for the Blue Crane in South Africa. The Ostrich.
56	Mewes, W., G. Nowald & H. Prange (2003): Kraniche - Mythen, Forschung, Fakten. Karlsruhe.
57	Morais, M., in litt. 2005.
58	Morozov, V.V. 2000. Current status of the southern subspecies of the Whimbrel <i>Numenius phaeopus alboaxillaris</i> (Lowe 1921) in Russia and Kazakstan. Wader Study Group Bulletin 92: 30-37.
59	Morrison, R.I.G., McCaffery, B.J., Skagen, S., Andres, B., Page, G., Jones, S., and Gill, R.E. 2005. Population estimates of North American shorebirds, 2005. Draft MS.
60	Nilsson, L., & Thomas Heinicke in litt 2005.
61	Overdijk, O. & Zwarts, F. 2002 (in prep). Proceedings of the fourth Eurosite Spoonbill workshop, Texel, The Netherlands, April 2002.
62	Overdijk, O., in litt. 2005.
63	Parasharya & Tere in litt. to SR, 2005. Unpublished Report: Flamingo Population Estimates in India, 1945-2003.
64	Parker, Vincent, in litt., 2002. Information provided for draft Atlas of Wader Populations in Africa and West Eurasia.
65	Perennou, C. 1991. Les recensements internationaux d'oiseaux d'eau en Afrique Tropicale. IWRB Spec. Publ. No. 15. Slimbridge, UK. 140 pp.
66	Perennou, C. 1992. African waterfowl census 1992. Les Dénombrements Internationaux d'oiseaux d'eau en Afrique, 1992. IWRB, Slimbridge, U.K. 87 pp.
67	PERSGA/GEF. 2003. Status of Breeding Seabirds in the Red Sea and Gulf of Aden. PERSGA Technical Series No. 8. PERSGA, Jeddah.
68	Pollitt, M., Hall, C. Holloway, S., Hearn, R., Marshall, P., Musgrove, A., Robinson, J., & Cranswick, P. 2003. The Wetland Bird Survey 2000-01: Wildfowl & Wader Counts. BTOWWT/RSPB/JNCC, Slimbridge, UK
69	van Roomen, M, van Winden, E., Koffijberg, K., Boele, A., Hustings, F., Kleefstra, R., Schoppers, J., van Turnhout, C., SOVON Ganzen en zwanenwerkgroep & Soldaat, L. 2004. Watervogels in Nederland in 2002/2003. SOVON -monitoringrapport 2004/02, RIZA-Rapport BM04/09, SOVON Vogelonderzoek, Nederland, Beek-Ubbergen.
70	Savchenko, A.P., Sidorkin, V.N. & Belyakov, A.V. 2001. Animal world of the Yenisei Plain. Vol.1. Amphibians, reptiles, birds. Krasnoyarsk, Krasnoyarsk State University. (In Russian.)
71	Schricke, V., & P. Triplet in litt. 2005.
72	Scott, D.A. and Rose, P.M. 1996. Atlas of Anatidae populations in Africa and western Eurasia. Wetlands International Publication No. 41. Wetlands International, Wageningen, NL. 336 pp.
73	Scott, D.A. in press, 2002. Report on the Conservation Status of Migratory Waterbirds in the Agreement Area. Update Report to African Eurasian Migratory Waterbird Agreement Secretariat
74	Simmons, R.E., Cordes, I. & Braby, R. 1998. Latitudinal trends, population size and habitat preferences of the Damara Tern <i>Sterna balaenarum</i> on Namibia's desert coast. Ibis 140: 439-445.
75	Specialist]] Unpublished information supplied by Wetlands International Specialist Groups.
76	Stroud, D.A., Davidson, N.C., West, R., Scott, D.A., Haanstra, L., Thorup, O., Ganter, B. & Delany, S. (compilers) on behalf of the International Wader Study Group 2004. The status of migratory wader populations in Africa and Western Eurasia in the 1990s. International Wader Studies 15: 259pp. http://web.uct.ac.za/depts/stats/adu/wsg/index.html
77	Syroechkovski, E.E., Jnr. 2002. Distribution and population estimates for swans in the Siberian arctic in the 1990s. In Proceedings of the Fourth International Swan Symposium, 2001 (ed. E.C. Rees, S.L. Earnst and J. Coulson). Waterbirds, Special Edition No. 1: 100-113.
78	Taylor, P.B. & van Perlo, B. (1998). Rails: a Guide to the Rails, Crakes, Gallinules and Coots of the World. Pica Press, Mountfield, Sussex.
79	Thévenot, M., Vernon, R. & Bergier, P. 2003. The Birds of Morocco: an annotated Checklist. BOU Checklist No. 20, Tring.
80	Thorup, O. 2004. Breeding Waders in Europe. International Wader Studies.

81	du Toit, M., Boere, G.C., Cooper, J., Kemper, J., Lenten, B., Simmons, R.S., Whittington, P.A. & Byers, O. (eds). 2002. Conservation assessment and management plan for southern African coastal seabirds. Workshop Report, Cape Town, South Africa, 4-8 February 2002. Avian Demography Unit, Cape Town & Conservation Breeding Specialist Group, IUCN/SSC
82	Torres Esquivias, J.A. & Andy Green in litt. June 2002.
83	Tree, A.J. & Klages, N.T.W. 2004. Population size, distribution and origins of Antarctic Terns <i>Sterna vittata</i> wintering in South Africa. <i>Marine Ornithology</i> 32: 55-61.
84	Tree, T., In litt. 2005.
88	Trolliet, B. & Girard, O. (2004). Anatid numbers and distribution in West Africa in winter. Abstracts of the Waterbirds Around the World Conference, Edinburgh, UK, April 2004, p. 299.
85	Trolliet, B. in litt.
87	Trolliet, B., & Fouquet, M. 2002. La population ouest-africaine du Flamant nain <i>Phoeniconias minor</i> : effectifs, répartition et isolement. <i>Malimbus</i> : 24 : 87-92.
89	Trolliet, B., & Fouquet, M. 2004. Wintering waders in coastal Guinea. <i>Wader Study Group Bull.</i> 103 : 56-62
86	Trolliet, B., & Girard, O., 2001. Numbers of Ruff <i>Philomachus pugnax</i> , wintering in West Africa. <i>Wader Study Group Bulletin</i> 96: 74-78.
90	Trolliet, B., Fouquet, M., & Keita, N. (in press, 2004). Statut du Flaman nain en Afrique de l'Ouest. <i>Proc. of the 11th Pan-African Ornithological Congress</i> .
91	Underhill, L., in litt. 2002.
92	Veen, J. Peeters, J., Leopold, M., van Damme, C., & Veen, P. In press, 2002. Effecten van visserij op natuurwaarden langs de kust van West Afrika: visetende vogels als graadmeters voor de kwaliteit van het marine milieu. Alterra rapport, wageningen, The Netherlands.
93	Veen, J., In litt. 2005.
94	Vidal, E. Duhem, C., Beau brun, P.C. & Yesou, P. (in press, 2002) Goeland leucophee Larus michahallis. In: Cadiou, B., Pons, J.M. & Yesou, P. (eds). Oiseaux marans nicheurs de France metropolitaine. Publication due Sept-Nov 2002.
95	Wetlands International, 2002. Waterbird Population Estimates - Third Edition. Wetlands International Global Series No. 12. Wageningen, The Netherlands.
96	Wetlands International: International Waterbird Census data.
97	Worden, J., Mitchell, C.R., Merne, O.J. & Cranswick, P.A. 2004. Greenland Barnacle Geese <i>Branta leucopsis</i> in Britain and Ireland: results of the international census, March 2003. Unpublished Report, WWT, Slimbridge.
98	WWT data.
99	WWT/JNCC Goose and Swan Monitoring Programme data.
100	Zydelis, R., Lorentsen, S.-H., Fox, A.D., Kuresoo, A., Bustnes, J.-O., Hario, M., Nilsson L. and Stipniece A. 2004. Recent changes in the status of Steller's Eider <i>Polysticta stelleri</i> wintering in Europe. (submitted for publication to <i>Bird Conservation International</i>).

TABLE 2: POPULATION SIZES AND TRENDS OF WATERBIRD SPECIES PROPOSED FOR INCLUSION IN THE AGREEMENT

Population	Estimate 2005	Trend 2005
<i>Phaethon aethereus</i> Red-billed Tropicbird		
<i>aethereus</i> - South Atlantic	<9,000	?
<i>indicus</i> - Persian Gulf, Gulf of Aden, Red Sea	(<1000)	?
<i>Phaethon rubricauda</i> Red-tailed Tropicbird		
<i>rubricauda</i> - Indian Ocean	(A)	?
<i>Phaethon lepturus</i> White-tailed Tropicbird		
<i>lepturus</i> - Persian Gulf, Gulf of Aden, Red Sea	7,500	?
<i>Sula (Morus) bassana</i> Northern Gannet	675,000	?
<i>Sula dactylatra</i> Masked Booby		
<i>melanops</i> – W Indian Ocean	(A)	DEC
<i>Fregata minor</i> Great Frigatebird		
<i>aldabrensis</i> - W Indian Ocean	(A)	?
<i>Fregata ariel</i> Lesser Frigatebird		
<i>iredalei</i> – W Indian Ocean	(A)	?
<i>Catharacta skua</i> Great Skua	40,800	?
<i>Stercorarius longicaudus</i> Long-tailed Skua		
<i>longicaudus</i>	(C/D)	?
<i>Rissa tridactyla</i> Black-legged Kittiwake		
<i>tridactyla</i>	E	?
<i>Sterna anaethetus</i> Bridled Tern		
<i>melanopterus</i> – W Africa	1,500	DEC
<i>fuligula</i> – Red Sea, E Africa, Persian Gulf, Arabian Sea to W India	390,000	
<i>antarctica</i> – S Indian Ocean	(C)	?
<i>Sterna fuscata</i> Sooty Tern		
<i>nubilosa</i> – Red Sea, Gulf of Aden, E to Pacific	(D/E)	?
<i>Anous stolidus</i> Brown Noddy		
<i>plumbeigularis</i> – Red Sea & Gulf of Aden	75,000	?
<i>Anous minutus</i> Black Noddy		
<i>atlanticus</i> – Atlantic Islands, Gulf of Guinea	120,000-180,000	?
<i>Anous tenuirostris</i> Lesser Noddy		
<i>tenuirostris</i> –Indian OceanIslands to E Africa	(B/C)	?
<i>Alle alle</i> Little Auk		
<i>alle</i> High Arctic, Baffin Is – Novaya Zemlya	36,000,000-	?

	39,000,000	
<i>Uria aalge</i> Common Guillemot		
<i>aalge</i> – E North America, Greenland, Iceland, Faeroes, Scotland, S Norway, Baltic	(D)	?
<i>albionis</i> Ireland, S Britain, France, Iberia, Helgoland	(D)	?
<i>hyperborea</i> Svalbard, N Norway to Novaya Zemlya	(D)	?
<i>Uria lomvia</i> Brunnich's Guillemot		
<i>lomvia</i> – E North America, Greenland, E to Severnaya Zemlya	9,000,000-12,000,000	?
<i>Alca torda</i> Razorbill		
<i>torda</i> E North America, Greenland, E to Baltic & White Seas	240,000	?
<i>islandica</i> Iceland, Faeroes, Britain, Ireland, Helgoland, NW France	1,950,000	?
<i>Cepphus grylle</i> Black Guillemot		
<i>grylle</i> Baltic Sea	(B/C)	?
<i>mandtii</i> Arctic E North America to Greenland, Jan Mayen & Svalbard E through Siberia to Alaska	(C)	?
<i>arcticus</i> N America, S Greenland, Britain, Ireland, Scandinavia, White Sea	(C)	?
<i>islandicus</i> Iceland	50,000	?
<i>faeroeensis</i> Faeroes	(B/C)	?
<i>Fratercula arctica</i> Atlantic Puffin		
<i>arctica</i> Hudson bay & Maine E to S Greenland, Iceland, Bear Is, Norway to S Novaya Zemlya	E	?
<i>naumanni</i> NE Canada, N Greenland, to Jan Mayen, Svalbard, N Novaya Zemlya	E	?
<i>grabae</i> Faeroes, S Norway & Sweden, Britain, Ireland, NW France	13,500,000	?